arXiv:1202.5052v1 [math-ph] 22 Feb 2012

Dyson's Brownian Motion Model as a Special Case of Dunkl Processes and Dunkl's Intertwining Operators

Sergio Andraus,[1,*] Makoto Katori,[2,†] and Seiji Miyashita[1,‡]

[1] *Department of Physics, Graduate School of Science, University of Tokyo*
7-3-1, Hongo, Bunkyo-ku, Tokyo, 113-0033

[2] *Department of Physics, Graduate School of Science and Engineering, Chuo University*
1-13-27 Kasuga, Bunkyo-ku, Tokyo 112-8551

Dyson's Brownian motion model is a one-parameter family of log-potential interacting particle systems in one dimension parametrized by an inverse temperature $\beta > 0$. When $\beta = 1, 2$ and 4, this model can be regarded as a stochastic realization of the eigenvalue statistics of Gaussian random matrices. Dunkl processes are mathematically defined using differential-difference operators (Dunkl operators) associated with finite abstract vector sets called root systems. When the root system is specified to be of type A, Dunkl processes constitute a one-parameter family of interacting particles in one dimension, in which particles perform not only diffusive motion and mutual repulsion but also interchange positions spontaneously. In the present paper, we prove that the type-A Dunkl processes with parameter $k > 0$ starting from any symmetric initial configuration are equivalent to Dyson's model with the inverse temperature $\beta = 2k$. We focus on the intertwining operators, since they play a central role in the mathematical theory of Dunkl operators, but their general closed form is not yet known. Using the equivalence between symmetric Dunkl processes and Dyson's model, we extract the effect of the A-type intertwining operator on symmetric polynomials from these processes' transition probability densities. In the zero-temperature limit, the intertwining operator maps all symmetric polynomials onto a function of the sum of their variables. This allows for the analysis of the zero-temperature limit of Dyson's model, which becomes a deterministic process with a final configuration proportional to a vector of the roots of the Hermite polynomials multiplied by the square root of the process time, while being independent of the initial configuration.

PACS numbers: 05.40.Jc, 02.50.-r, 02.30.Gp

---

[*] andraus@spin.phys.s.u-tokyo.ac.jp

[†] katori@phys.chuo-u.ac.jp

[‡] miya@spin.phys.s.u-tokyo.ac.jp

I. INTRODUCTION

Dyson's Brownian motion model was introduced as the stochastic process that the eigenvalues of a Gaussian random matrix undergo when its entries are independent Brownian motions [1]. In this sense, Dyson's model is a dynamic version of Gaussian random matrices [2], and therefore both models share the same parameter β for each of the three Gaussian ensembles of random matrices, 1, 2 and 4 for the Gaussian orthogonal, unitary and symplectic ensembles respectively [2–4]. While β is usually considered as a discrete parameter, in the case of Dyson's model it can be regarded as a real positive parameter. In that case, Dyson's model is a system of N Brownian particles in one dimension that repel each other via a log-potential, and $\beta/2$ is its coupling constant [5]. Additionally, in view of the fact that the joint probability density function of the eigenvalues of Gaussian random matrices has the form of the partition function of a one-dimensional log-gas [3], β also carries the meaning of inverse temperature. The vicious walker model formulated by Fisher [6] consists of N random walkers who annihilate each other if they make contact; one of the most particular properties of Dyson's model is that for $\beta = 2$, Dyson's model is a diffusion scaling limit of the vicious walker model with the restriction that all walkers must survive at all times [7]. This restriction is equivalent to conditioning the N particles never to collide, and therefore this process is also known as the non-colliding Brownian motion [8, 9]. Due to this connection, Dyson's model has found applications in polymer networks [10, 11], traffic models [12], gauge fields and nuclear physics [13–17], and random growth models [18, 19], among others. In particular, Dyson's model has been applied extensively in the Kardar-Parisi-Zhang universality class [20–23], and there have been recent experimental observations of phenomena pertaining to it that confirm theoretical predictions [24, 25]. Also, because of this relationship, Dyson's model is intimately related to the mathematical areas of combinatorics, group theory and representation theory [26–29]. Therefore, Dyson's model has been an active topic of research since its introduction, and remains one of the most important models in non-equilibrium statistical mechanics and in stochastic process theory as well as a topic of great interest in mathematics.

On the other hand, Dunkl operators are differential-difference operators [30] that depend on a series of parameters and a finite set of vectors. This set of vectors is called root system. In this paper, we will focus on the A-type root system and its corresponding Dunkl operators given by [31]

$$T_i f(\boldsymbol{x}) = \frac{\partial}{\partial x_i} f(\boldsymbol{x}) + k \sum_{\substack{j:j=1\\ j\neq i}}^{N} \frac{f(\boldsymbol{x}) - f(\sigma_{ij}\boldsymbol{x})}{x_i - x_j}, \tag{1}$$

where $k \geq 0$ is a real scalar parameter, $\boldsymbol{x}$ is a vector in $\mathbb{R}^N$, $\sigma_{ij}\boldsymbol{x}$ denotes the vector $\boldsymbol{x}$ with its i-th and j-th components interchanged and the function f is an arbitrary differentiable scalar function. This and other types of Dunkl operators have been used in mathematical physics in the study of integrable systems and quantum many-body models [32–34], but the examples of applications of these operators are scarce in comparison to Dyson's model. While initially defined as a tool to study multivariate orthogonal polynomials [30], Dunkl operators have also been used in harmonic analysis [35] and stochastic processes [36].

Dunkl processes are defined as follows [37]. Consider first the N-dimensional heat equation. The N-dimensional heat kernel is the transition probability density (TPD) that corresponds to the heat equation as a Kolmogorov backward equation. Therefore, one can use the heat kernel to define N-dimensional Brownian motion. Now, replacing the spatial partial derivatives in the heat equation with Dunkl operators such as the one given in Eq. (1) yields the Dunkl heat equation. Then, the TPD that solves the Dunkl heat equation as a Kolmogorov backward equation defines a generalization of Brownian motion. Dunkl processes are defined as this generalization, that is, as the family of stochastic processes that obey the TPD that corresponds to the Dunkl heat equation. Hence, different root systems define different kinds of Dunkl processes. In the case of the type-A Dunkl operators (1), the corresponding A-type Dunkl processes are a one-parameter family of stochastic processes, where the real parameter k is non-negative [37]. Dunkl processes subject to symmetric initial conditions with respect to their respective root systems are called radial Dunkl processes [38, 39], and their properties have been an active topic of research in recent years [40, 41]. In order to emphasize their variable exchange symmetry, we will refer to the A-type radial Dunkl processes as symmetric Dunkl processes.

Many of the results from the theory of Dunkl operators are achieved through the use of Dunkl's intertwining operator V_k [42], which we review in Section III. Consider a function $f(\boldsymbol{x})$ as in Eq. (1). Then, V_k is defined by the following equation:

$$T_i V_k f(\boldsymbol{x}) = V_k \frac{\partial}{\partial x_i} f(\boldsymbol{x}). \tag{2}$$

In other words, V_k binds Dunkl operators with regular partial derivatives. Additionally, it has no effect on constants (i.e., $V_k 1 = 1$) and when it is applied on a monomial of a given degree, it produces a homogeneous polynomial of the same total degree. Note that there is a different operator for each type of Dunkl operator. Through the use of V_k, many calculations are simplified, as it allows Dunkl operators to be treated as partial derivatives. However, it presently has no general closed form.

After reviewing Dyson's model briefly in Sec. II A, we will show in Theorem 1, Sec. II B, that the TPD of symmetric Dunkl processes obey a Kolmogorov backward equation that is identical to that of Dyson's model. Therefore, their TPDs are equivalent and a relationship between the Dunkl parameter k and the parameter β in Dyson's model is established. This relationship provides a physical meaning to the abstract Dunkl parameter k, that is, k can be understood as a parameter proportional to the inverse temperature. In Sec. III, we will review the intertwining operator and its defining properties. Using Theorem 1, we will extract the effect of the intertwining operator on symmetric polynomials from the TPDs of Dyson's model and symmetric Dunkl processes. This is the result of Theorem 2, Section IV A. In order to gain a better understanding of this result, we will make observations on the particular case of quadratic symmetric polynomials in Section IV B. We will find that in the $k \to \infty$ limit, V_k turns all symmetric functions into a function of the sum of their variables. We will prove this fact in Theorem 3, Section IV C. Finally, we will make use of the behavior of $V_{k\to\infty}$ to investigate the limit $k = \beta/2 \to \infty$ of the TPD of the symmetric Dunkl process (or its equivalent Dyson's model) in Theorem 4, Section V. We will find that in this limit, this process becomes deterministic and its final configuration is proportional to the roots of the Hermite polynomials multiplied by the square root of the process time. We will also find that the form of $V_{k\to\infty}$ is responsible for making the symmetric Dunkl process independent of its initial condition in the zero-temperature limit.

II. DYSON'S BROWNIAN MOTION MODEL AND DUNKL PROCESSES

A. Dyson's Brownian Motion Model

Dyson's Brownian motion model [1] is a stochastic process in which N particles undergo Brownian motion in one dimension while interacting repulsively with each other with a strength proportional to the inverse of the distances between them. Let us denote the Laplacian operator by

$$\Delta^{(x)} = \sum_{i=1}^{N} \frac{\partial^2}{\partial x_i^2}. \tag{3}$$

Dyson's model is described by the Kolmogorov backward equation (see, e.g., Remark 1 in Ref. [5])

$$\frac{\partial}{\partial t} P_\beta(t, \boldsymbol{y}|\boldsymbol{x}) = \frac{1}{2}\Delta^{(x)} P_\beta(t, \boldsymbol{y}|\boldsymbol{x}) + \frac{\beta}{2} \sum_{i=1}^{N} \sum_{\substack{j=1: \\ j\neq i}}^{N} \frac{1}{x_i - x_j} \frac{\partial}{\partial x_i} P_\beta(t, \boldsymbol{y}|\boldsymbol{x}), \tag{4}$$

where $P_\beta(t, \boldsymbol{y}|\boldsymbol{x})$ is the TPD that the particles reach the positions $\boldsymbol{y} = (y_1, \ldots, y_N)$ after a time t, given that they started from $\boldsymbol{x} = (x_1, \ldots, x_N)$. While we denote vectors in $\mathbb{R}^N$ using boldface symbols, we will use the notation $x^2 = \boldsymbol{x} \cdot \boldsymbol{x}$ to denote their squared norm.

The eigenvalues $\boldsymbol{\zeta} = \{\zeta_i\}_{i=1,\ldots,N}$ of the Gaussian ensembles of random matrices obey the joint probability density function (Theorem 3.3.1 of Ref. [3])

$$\mathrm{Prob}_{N,\beta}(\boldsymbol{\zeta}) \propto \exp(-\beta\mathcal{H}) \tag{5}$$

with

$$\mathcal{H} = \frac{1}{2}\zeta^2 - \sum_{1\le i<j\le N} \log|\zeta_j - \zeta_i|. \tag{6}$$

Because $\mathrm{Prob}_{N,\beta}(\boldsymbol{x})$ has the form of the partition function of a log-gas in a harmonic potential in one dimension after integrating out the kinetic part, one can consider β as a real positive parameter by giving it the meaning of inverse temperature in this log-gas model. In the same way, one can treat β as a positive real number in Eq. (4) by regarding $\beta/2$ as the coupling constant of the interaction (drift) term. Therefore, when β is extended from the three values 1,2 and 4, it carries two different physical meanings. Loosely speaking, the inverse-temperature interpretation of β still holds in Dyson's model: when $\beta \to 0$, the interaction term in Eq. (4) vanishes leaving an N-dimensional Brownian motion, which means that the fluctuations in the system are much larger in magnitude than the interaction between particles. This is the analog of a high temperature condition. On the other hand, when $\beta \to \infty$, the diffusion term is negligible compared to the interaction term, and so the randomness in the system disappears, which is the analog of a zero-temperature condition.

A consequence of the results in Refs. [43] and [36] (see, e.g., Ref. [38]) is that, for N-dimensional vectors $\boldsymbol{x}$ and $\boldsymbol{y}$ ordered so that $y_i < y_j$ and $x_i < x_j$ for $i < j$, the TPD of Dyson's model is given by

$$P_\beta(t, \boldsymbol{y}|\boldsymbol{x}) = \frac{N! e^{-(x^2+y^2)/2t}}{(2\pi t)^{N/2}} \prod_{j=1}^{N} \left[\frac{\Gamma(1+\beta/2)}{\Gamma(1+j\beta/2)}\right] \left|h_N\left(\frac{\boldsymbol{y}}{\sqrt{t}}\right)\right|^\beta {}_0\mathcal{F}_0^{(2/\beta)}\left(\frac{\boldsymbol{x}}{\sqrt{t}}, \frac{\boldsymbol{y}}{\sqrt{t}}\right), \tag{7}$$

where $\Gamma(x)$ is the gamma function, $h_N(\boldsymbol{y}) = \prod_{1\le i<j\le N}(y_j - y_i)$ is the Vandermonde determinant and ${}_0\mathcal{F}_0^{(2/\beta)}$ is the generalized hypergeometric function (see Appendix A). In the case $\beta = 2$, the non-colliding Brownian motion, Dyson's model has the following TPD as shown by Grabiner [8]:

$$P_{\beta=2}(t, \boldsymbol{y}|\boldsymbol{x}) = \frac{h_N(\boldsymbol{y})}{h_N(\boldsymbol{x})} \det_{1\le i,j\le N} \left(\frac{e^{-(x_i-y_j)^2/2t}}{\sqrt{2\pi t}}\right). \tag{8}$$

B. Dunkl Processes

Within our present setting, we are interested in the A-type Dunkl operators given by Eq. (1) (see Appendices B and C). Using these operators, one can define Dunkl processes as follows [36]: consider the TPD, $p_k(t,\boldsymbol{y}|\boldsymbol{x})$, that solves the Dunkl heat equation

$$\frac{\partial}{\partial t}p_k(t,\boldsymbol{y}|\boldsymbol{x}) = \frac{1}{2}\sum_{i=1}^{N} T_i^2 p_k(t,\boldsymbol{y}|\boldsymbol{x}) \tag{9}$$

as its Kolmogorov backward equation. Then the stochastic process that obeys $p_k(t,\boldsymbol{y}|\boldsymbol{x})$ as its TPD is defined as the Dunkl process of type A. Direct insertion of Eq. (1) into Eq. (9) yields

$$\begin{aligned}\frac{\partial}{\partial t}p_k(t,\boldsymbol{y}|\boldsymbol{x}) = {} & \frac{1}{2}\Delta^{(x)}p_k(t,\boldsymbol{y}|\boldsymbol{x}) + \sum_{i=1}^{N}\sum_{\substack{j=1:\\ j\neq i}}^{N}\frac{k}{x_i-x_j}\frac{\partial}{\partial x_i}p_k(t,\boldsymbol{y}|\boldsymbol{x}) \\ & -\frac{k}{2}\sum_{i=1}^{N}\sum_{\substack{j=1:\\ j\neq i}}^{N}\frac{p_k(t,\boldsymbol{y}|\boldsymbol{x})-p_k(t,\boldsymbol{y}|\sigma_{ji}\boldsymbol{x})}{(x_j-x_i)^2}.\end{aligned} \tag{10}$$

Reading off the terms of this Kolmogorov backward equation gives us information about Dunkl processes: the first term, a Laplacian acting on $\boldsymbol{x}$, is a diffusion term, while the second term is a drift term that drives the process away from the surfaces defined by the equations $x_i = x_j$ for any $i \neq j$. The third term is a spontaneous jump term, which instantaneously makes the process jump from $\boldsymbol{x}$ to some permutation of itself so long as $p_k(t,\boldsymbol{y}|\boldsymbol{x})$ is not symmetric under permutations of $\boldsymbol{x}$.

With the observation about the jump term in mind, the comparison of Eqs. (4) and (10) reveals a clear similarity between the two Kolmogorov backward equations. Let us define the symmetric group S_N as the set of all the permutations of N objects with composition as its group operation. The permutation $\rho \in S_N$ of the vector $\boldsymbol{x}$ is given by

$$\rho\boldsymbol{x} = (x_{\rho(1)},\ldots,x_{\rho(N)}). \tag{11}$$

We will consider the TPD of a Dunkl process with a symmetric initial condition. We will refer to such a process as a symmetric Dunkl process. Given the symmetric distribution

$$\mu^s_{\boldsymbol{x}}(\boldsymbol{z}) = \sum_{\rho\in S_N}\delta^N(\boldsymbol{z}-\rho\boldsymbol{x}), \tag{12}$$

where we define $\delta^N(\boldsymbol{x}-\boldsymbol{y}) = \prod_{j=1}^{N}\delta(x_j-y_j)$, we write

$$p^s_k(t,\boldsymbol{y}|\boldsymbol{x}) = \int_{\mathbb{R}^N} p_k(t,\boldsymbol{y}|\boldsymbol{x})\mu^s_{\boldsymbol{x}}(\boldsymbol{z})\,\mathrm{d}^N z = \sum_{\rho\in S_N} p_k(t,\boldsymbol{y}|\rho\boldsymbol{x}), \tag{13}$$

with $\mathrm{d}^N z = \prod_{j=1}^N \mathrm{d}z_j$. We show that the Kolmogorov backward equation obeyed by the TPD of the symmetric Dunkl process, $p_k^s(t, \boldsymbol{y}|\boldsymbol{x})$, is identical to that of the TPD of Dyson's model in the following theorem.

Theorem 1. *The TPD of a symmetric Dunkl process, p_k^s, solves the Kolmogorov backward equation of Dyson's model with parameter*

$$\beta = 2k. \tag{14}$$

That is, the equivalence

$$p_k^s(t, \boldsymbol{y}|\boldsymbol{x}) = P_{\beta=2k}(t, \boldsymbol{y}|\boldsymbol{x}) \tag{15}$$

is established.

Proof. Comparing Eqs. (10) and (4), we see that the only difference between the two is the third term on the RHS of Eq. (10). The bijective nature of permutations allows us to write

$$\Delta^{(x)} = \sum_{i=1}^N \frac{\partial^2}{\partial x_{\rho(i)}^2},$$

$$\sum_{i=1}^N \sum_{\substack{j=1 \\ j\neq i}}^N \frac{1}{x_i - x_j} \frac{\partial}{\partial x_i} = \sum_{i=1}^N \sum_{\substack{j=1 \\ j\neq i}}^N \frac{1}{x_{\rho(i)} - x_{\rho(j)}} \frac{\partial}{\partial x_{\rho(i)}}$$

and

$$\sum_{i=1}^N \sum_{\substack{j=1 \\ j\neq i}}^N \frac{1-\sigma_{ij}}{(x_i - x_j)^2} = \sum_{i=1}^N \sum_{\substack{j=1 \\ j\neq i}}^N \frac{1-\sigma_{\rho(i)\rho(j)}}{(x_{\rho(i)} - x_{\rho(j)})^2}$$

for any permutation $\rho \in S_N$. Therefore,

$$\begin{aligned}
&\left[\frac{1}{2}\Delta^{(x)} + \sum_{i=1}^N \sum_{\substack{j=1 \\ j\neq i}}^N \frac{k}{x_i - x_j} \frac{\partial}{\partial x_i} - \frac{k}{2} \sum_{i=1}^N \sum_{\substack{j=1 \\ j\neq i}}^N \frac{1-\sigma_{ij}}{(x_i - x_j)^2}\right] \sum_{\rho\in S_N} p_k(t, \boldsymbol{y}|\rho\boldsymbol{x}) \\
&= \sum_{\rho\in S_N} \left[\frac{1}{2}\sum_{i=1}^N \frac{\partial^2}{\partial x_{\rho(i)}^2} + \sum_{i=1}^N \sum_{\substack{j=1 \\ j\neq i}}^N \frac{k}{x_{\rho(i)} - x_{\rho(j)}} \frac{\partial}{\partial x_{\rho(i)}} - \frac{k}{2} \sum_{i=1}^N \sum_{\substack{j=1 \\ j\neq i}}^N \frac{1-\sigma_{\rho(i)\rho(j)}}{(x_{\rho(i)} - x_{\rho(j)})^2}\right] p_k(t, \boldsymbol{y}|\rho\boldsymbol{x}) \\
&= \sum_{\rho\in S_N} \frac{\partial}{\partial t} p_k(t, \boldsymbol{y}|\rho\boldsymbol{x}) = \frac{\partial}{\partial t} p_k^s(t, \boldsymbol{y}|\boldsymbol{x}),
\end{aligned} \tag{16}$$

and in particular, since $p_k^s(t, \boldsymbol{y}|\boldsymbol{x})$ is symmetric in $\boldsymbol{x}$, the third term in the first line vanishes, leaving us with Eq. (4) provided we set $k = \beta/2$. □

Note that, because $p_k(t,\boldsymbol{y}|\boldsymbol{x})$ is a probability density function on $\boldsymbol{y}$, its integral with respect to $\boldsymbol{y}$ over $\mathbb{R}^N$ is equal to one, and so $p_k^s(t,\boldsymbol{y}|\boldsymbol{x})$ integrates to $N!$. A consequence of this theorem is that, because both P_β and p_k^s follow the same Kolmogrov backward equation, they must be equivalent. We will exploit this fact in Section IV.

III. THE INTERTWINING OPERATOR

In the previous section we obtained some information about the TPD of Dunkl processes and how it is related to Dyson's model. However, this TPD can also be calculated using the intertwining operator V_k, Eq. (2) [31, 42] (see Appendix D for details). While we will treat V_k from a mostly general perspective in this section, we will assume that V_k is the A-type intertwining operator.

Dunkl originally defined V_k for the case in which the function $f(\boldsymbol{x})$ in Eq. (2) is a polynomial, and proved its existence for $k \geq 0$ using a recursive construction. However, its explicit form is unknown in general, and known only in a few particular cases. From Eq. (1), one can note that when $k = 0$, the intertwining operator is the identity operator.

The intertwining operator relates N-dimensional Brownian motion to Dunkl processes as follows. Let us consider the N-dimensional heat equation

$$\left(\frac{\partial}{\partial t} - \frac{1}{2}\Delta^{(x)}\right) p_0(t,\boldsymbol{y}|\boldsymbol{x}) = 0 \tag{17}$$

with its solution

$$p_0(t,\boldsymbol{y}|\boldsymbol{x}) = \frac{e^{-(x^2+y^2)/2t}}{(2\pi t)^{N/2}} e^{\boldsymbol{x}\cdot\boldsymbol{y}/t} \tag{18}$$

given the initial condition $p_0(0,\boldsymbol{y}|\boldsymbol{x}) = \delta^N(\boldsymbol{x}-\boldsymbol{y})$. This solution is both the Green function of the diffusion equation and the TPD of the N-dimensional Brownian motion. When we apply V_k on this equation, we obtain

$$0 = V_k\left(\frac{\partial}{\partial t} - \frac{1}{2}\Delta^{(x)}\right) p_0(t,\boldsymbol{y}|\boldsymbol{x}) = \left(\frac{\partial}{\partial t} - \frac{1}{2}\sum_{i=1}^N T_i^2\right) V_k p_0(t,\boldsymbol{y}|\boldsymbol{x}), \tag{19}$$

because of Eq. (2) and the fact that V_k commutes with the partial derivative relative to time. From the above, we deduce that the Green function for the Dunkl heat equation, which we denote by $\Gamma_k(t,\boldsymbol{y}|\boldsymbol{x})$, is given by

$$\Gamma_k(t,\boldsymbol{y}|\boldsymbol{x}) = V_k p_0(t,\boldsymbol{y}|\boldsymbol{x}). \tag{20}$$

On the other hand, the Green function for the Dunkl heat equation can be calculated directly using the Dunkl transform [35], a generalization of the Fourier transform, in the same way that the

Fourier transform can be used to derive Eq. (18), see Ref. [36]. In order to define the Dunkl transform, one needs to calculate the eigenfunction ϕ_k of the Dunkl operators $\{T_i\}_{i=1,\dots,N}$ of parameter k with eigenvalue $\boldsymbol{\xi}$,

$$T_i\phi_k(\boldsymbol{x},\boldsymbol{\xi}) = \xi_i\phi_k(\boldsymbol{x},\boldsymbol{\xi}), \quad i = 1, 2, \dots, N, \tag{21}$$

subject to $\phi_k(\boldsymbol{0},\boldsymbol{\xi}) = 1$ with $\boldsymbol{0} = (0, \dots, 0)$. When $k = 0$, Eq. (21) is

$$\frac{\partial}{\partial x_i}\phi_0(\boldsymbol{x},\boldsymbol{\xi}) = \xi_i\phi_0(\boldsymbol{x},\boldsymbol{\xi}),$$

and its solution is

$$\phi_0(\boldsymbol{x},\boldsymbol{\xi}) = e^{\boldsymbol{\xi}\cdot\boldsymbol{x}}. \tag{22}$$

This eigenfunction is used as the kernel of the N-dimensional Fourier transform with $\boldsymbol{x}$ replaced by $\sqrt{-1}\boldsymbol{x}$. With Eq. (22) in mind, we adopt the form

$$\phi_k(\boldsymbol{x},\boldsymbol{\xi}) = V_k e^{\boldsymbol{\xi}\cdot\boldsymbol{x}}. \tag{23}$$

Insertion of Eq. (23) into Eq. (21) yields

$$T_i\phi_k(\boldsymbol{x},\boldsymbol{\xi}) = T_iV_ke^{\boldsymbol{\xi}\cdot\boldsymbol{x}} = V_k\frac{\partial}{\partial x_i}e^{\boldsymbol{\xi}\cdot\boldsymbol{x}} = V_k\xi_ie^{\boldsymbol{\xi}\cdot\boldsymbol{x}} = \xi_iV_ke^{\boldsymbol{\xi}\cdot\boldsymbol{x}} = \xi_i\phi_k(\boldsymbol{x},\boldsymbol{\xi}), \tag{24}$$

where the second equality follows from the definition of the intertwining operator and the fourth equality follows from the fact that V_k is linear. In addition, when $\phi_k(\boldsymbol{x},\boldsymbol{\xi})$ is written as a power series, each term is a homogeneous polynomial of $\boldsymbol{x}$, whose degree is conserved by V_k, and when the limit $\boldsymbol{x} \to \boldsymbol{0}$ is taken, all homogeneous polynomials of degree larger than zero vanish; the only remaining term (the zero-order polynomial) is equal to one. Therefore, $\phi_k(\boldsymbol{x},\boldsymbol{\xi})$ as given in Eq. (23) is the solution of Eq. (21) with the condition $\phi_k(\boldsymbol{0},\boldsymbol{\xi}) = 1$.

In analogy to Eq. (22), $\phi_k(\boldsymbol{x},\boldsymbol{\xi})$ has the following properties [44, 45]: given $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}$,

$$\begin{gathered}
\phi_k(\boldsymbol{x},\boldsymbol{y}) > 0, \\
|\phi_k(\sqrt{-1}\boldsymbol{x},\boldsymbol{y})| \leq 1, \\
\phi_k(\boldsymbol{x},\boldsymbol{y}) = \phi_k(\boldsymbol{y},\boldsymbol{x}), \\
\phi_k(\lambda\boldsymbol{x},\boldsymbol{y}) = \phi_k(\boldsymbol{x},\lambda\boldsymbol{y}), \quad \lambda \in \mathbb{C},
\end{gathered} \tag{25}$$

and

$$\phi_k(\rho\boldsymbol{x},\rho\boldsymbol{y}) = \phi_k(\boldsymbol{x},\boldsymbol{y}), \tag{26}$$

with $\rho \in S_N$. When the Dunkl operators are not of the A type, $\phi_k(\boldsymbol{x}, \boldsymbol{y})$ obeys property (26), but the operator ρ is given by the root system under consideration and Eq. (B1), and it is not an element of S_N in general (see Ref. [37]). Due to these properties, $\phi_k(\boldsymbol{x}, \boldsymbol{\xi})$ can be used as the kernel for a transformation similar to the Fourier transform by replacing $\boldsymbol{x}$ by $\sqrt{-1}\boldsymbol{x}$. For this reason, hereafter we denote this eigenfunction by

$$E_k(\boldsymbol{x}, \boldsymbol{\xi}) \equiv \phi_k(\boldsymbol{x}, \boldsymbol{\xi}). \tag{27}$$

This function is known as the Dunkl kernel [42].

Using the Dunkl transform mentioned above, one can calculate $\Gamma_k(t, \boldsymbol{y}|\boldsymbol{x})$ and the TPD of Dunkl processes in general (see Appendix D, Eq. (D24) for the explicit relationship between Γ_k and p_k). However, because the Dunkl kernel is contained in it, and the intertwining operator is unknown in general, this TPD is not explicitly specified. In our case, $p_k(t, \boldsymbol{y}|\boldsymbol{x})$ is given by

$$p_k(t, \boldsymbol{y}|\boldsymbol{x}) = \prod_{j=1}^{N} \frac{\Gamma(1+k)}{\Gamma(1+jk)} \left| h_N \left(\frac{\boldsymbol{y}}{\sqrt{t}} \right) \right|^{2k} \frac{e^{-(x^2+y^2)/2t}}{(2\pi t)^{N/2}} E_k \left(\frac{\boldsymbol{x}}{\sqrt{t}}, \frac{\boldsymbol{y}}{\sqrt{t}} \right). \tag{28}$$

The last factor on the right is the only unknown part of this TPD, as mentioned before. Let us note that in order to obtain p_k, we need the Dunkl kernel, which in turn requires us to calculate the effect of the intertwining operator on the exponential $e^{\boldsymbol{x}\cdot\boldsymbol{y}}$. Conversely, if we have p_k, we can obtain the Dunkl kernel, and by expanding it into sums of suitably chosen polynomials, we can extract the effect of the intertwining operator on any polynomial.

IV. EXTRACTION OF V_k FROM DYSON'S MODEL

A. Theorem

Here, we will obtain an expression for the effect of V_k on symmetric polynomials. Using Theorem 1, we combine Eqs. (13) and (28), equate the result to Eq. (7), and rescale $\boldsymbol{x}$ and $\boldsymbol{y}$ by a factor of $\sqrt{t}$ to obtain

$$\sum_{\rho \in S_N} E_k(\rho \boldsymbol{x}, \boldsymbol{y}) = N! {}_0\mathcal{F}_0^{(1/k)}(\boldsymbol{x}, \boldsymbol{y}). \tag{29}$$

From Eq. (29), we extract the effect of the intertwining operator when applied to a symmetric polynomial. As preparation, let us introduce the following definitions: consider an arbitrary integer partition, μ, as defined in Appendix A. Let l_j^μ represent the multiplicity of the j-th (distinct) part of μ, where the subscript P in l_P^μ refers to the number of distinct parts of μ. In the cases where

$l(\mu) < N$, there are $N - l(\mu)$ zero parts in μ and therefore $l^\mu_P = N - l(\mu)$ accounts for the multiplicity of zero parts in the first N parts of μ. For example, if $N = 6$ and $\mu = (5,3,2,2) = (5,3,2,2,0,0)$, then $P = 4$ and $l^\mu_1 = 1$, $l^\mu_2 = 1$, $l^\mu_3 = 2$ and $l^\mu_4 = 2$. Using this notation we define the following multinomial coefficient:

$$M(\mu, N) = \frac{N!}{l^\mu_1! \cdots l^\mu_P!}. \tag{30}$$

This function represents the number of distinct permutations of μ when it is considered as an N-dimensional vector. See Appendix A (in particular, Eqs. (A3), (A11), (A16) and (A17)) for the definitions of the monomial symmetric functions $m_\lambda(\boldsymbol{x})$, Jack functions $\mathcal{P}^{(1/k)}_\lambda(\boldsymbol{x})$ and the various quantities involved in the theorem below.

Theorem 2. *The action of the intertwining operator on a symmetric polynomial is given by the equation*

$$V_k m_\lambda(\boldsymbol{x}) = \lambda! M(\lambda, N) \sum_{\substack{\tau : l(\tau) \le N \\ |\tau| = |\lambda|}} \frac{c_\tau(1/k)}{c'_\tau(1/k)} \frac{u_{\tau\lambda}(1/k)}{(kN)^{(1/k)}_\tau} \mathcal{P}^{(1/k)}_\tau(\boldsymbol{x}). \tag{31}$$

That is, the intertwining operator of type A maps the monomial symmetric functions $m_\lambda(\boldsymbol{x})$ onto a linear combination of Jack functions of parameter $1/k$, $\mathcal{P}^{(1/k)}_\tau(\boldsymbol{x})$.

Proof. We first expand the LHS of Eq. (29) in terms of symmetric polynomials. For this purpose, we expand the symmetrized exponential $\sum_{\rho \in S_N} \exp(\rho\boldsymbol{x} \cdot \boldsymbol{y})$ in terms of the monomial symmetric functions $m_\mu(\boldsymbol{x})$.

$$\begin{aligned}
\sum_{\rho \in S_N} e^{\rho\boldsymbol{x}\cdot\boldsymbol{y}} &= \sum_{\rho \in S_N} \sum_{n=0}^{\infty} \sum_{\substack{\mu : l(\mu) \le N \\ |\mu| = n}} \frac{1}{\mu!} \sum_{\substack{\tau \in S_N : \\ \tau(\mu) \text{ distinct}}} \prod_{j=1}^{N} (x_{\rho(j)} y_j)^{\mu_{\tau(j)}} \\
&= \sum_{\mu : l(\mu) \le N} \frac{1}{\mu!} \sum_{\substack{\tau \in S_N : \\ \tau(\mu) \text{ distinct}}} \left\{ \sum_{\rho \in S_N} \prod_{j=1}^{N} x_{\rho(j)}^{\mu_{\tau(j)}} \right\} \prod_{j=1}^{N} y_j^{\mu_{\tau(j)}} \\
&= \sum_{\mu : l(\mu) \le N} \frac{1}{\mu!} \left\{ \sum_{\rho' \in S_N} \prod_{j'=1}^{N} x_{j'}^{\mu_{\rho'(j')}} \right\} \sum_{\substack{\tau \in S_N : \\ \tau(\mu) \text{ distinct}}} \prod_{j=1}^{N} y_j^{\mu_{\tau(j)}}
\end{aligned} \tag{32}$$

In the last line we have used the substitutions $j' = \rho(j)$ and $\rho'(j') = \tau[\rho^{-1}(j')]$. The last term on the right is, by definition, $m_\mu(\boldsymbol{y})$. The term inside the braces is equal to $m_\mu(\boldsymbol{x})$ multiplied by the number of non-distinct permutations of μ. Using Eq. (30), we write the above as

$$\sum_{\rho \in S_N} e^{\rho\boldsymbol{x}\cdot\boldsymbol{y}} = \sum_{\mu : l(\mu) \le N} \frac{N! m_\mu(\boldsymbol{x}) m_\mu(\boldsymbol{y})}{\mu! M(\mu, N)}. \tag{33}$$

Applying V_k on the above yields the LHS of Eq. (29) expanded in terms of a sum of monomial symmetric functions. The next step is to eliminate the variable $\boldsymbol{y}$ using the orthogonality of Jack functions. Insertion of Eq. (A12) in Eq. (33) after applying V_k yields

$$\sum_{\rho\in S_N} E_k(\rho\boldsymbol{x},\boldsymbol{y}) = V_k \sum_{\mu:l(\mu)\leq N} \frac{N! m_\mu(\boldsymbol{x})}{\mu! M(\mu,N)} \sum_{\substack{\nu:\nu\leq\mu\\ |\nu|=|\mu|}} (u^{-1})_{\mu\nu}(1/k)\mathcal{P}_\nu^{(1/k)}(\boldsymbol{y}). \tag{34}$$

Using Eqs. (34) and (A19), Eq. (29) becomes

$$V_k \sum_{\mu:l(\mu)\leq N} \frac{m_\mu(\boldsymbol{x})}{\mu! M(\mu,N)} \sum_{\substack{\nu:\nu\leq\mu\\ |\nu|=|\mu|}} (u^{-1})_{\mu\nu}(1/k)\mathcal{P}_\nu^{(1/k)}(\boldsymbol{y}) = \sum_{\tau:l(\tau)\leq N} \frac{c_\tau(1/k)}{c'_\tau(1/k)} \frac{\mathcal{P}_\tau^{(1/k)}(\boldsymbol{x})\mathcal{P}_\tau^{(1/k)}(\boldsymbol{y})}{(kN)_\tau^{(1/k)}}, \tag{35}$$

and from the orthogonality of Jack functions [29] and the linearity of V_k, which acts only on $\boldsymbol{x}$, we can equate the coefficients of the same Jack functions of $\boldsymbol{y}$ to obtain

$$\sum_{\substack{\mu:l(\mu)\leq N\\ |\mu|=|\tau|}} \frac{(u^{-1})_{\mu\tau}(1/k)}{\mu! M(\mu,N)} V_k m_\mu(\boldsymbol{x}) = \frac{c_\tau(1/k)}{c'_\tau(1/k)} \frac{\mathcal{P}_\tau^{(1/k)}(\boldsymbol{x})}{(kN)_\tau^{(1/k)}}. \tag{36}$$

This relation is solved for $V_k m_\lambda(\boldsymbol{x})$ if we apply the sum $\sum_\tau u_{\tau\lambda}(1/k)$, which yields

$$\frac{V_k m_\lambda(\boldsymbol{x})}{\lambda! M(\lambda,N)} = \sum_{\substack{\tau:l(\tau)\leq N\\ |\tau|=|\lambda|}} u_{\tau\lambda}(1/k) \frac{c_\tau(1/k)}{c'_\tau(1/k)} \frac{\mathcal{P}_\tau^{(1/k)}(\boldsymbol{x})}{(kN)_\tau^{(1/k)}}. \tag{37}$$

From this last operation, the theorem follows. □

From Theorem 2, we can recover known results from the cases $k = 0$ (where V_k must reduce to the identity operator), $k = 1$ (which should yield the known TPD of the non-colliding Brownian motion) and the case in which all particles start from the same position. We rederive these results in detail in Appendix E.

B. Effect of V_k on symmetric quadratic equations

Although Theorem 2 is consistent with known results, it does not give an immediate understanding of the effect of V_k on symmetric polynomials. The purpose of this subsection is to consider a simple case of Theorem 2 and make observations of the general properties of V_k in order to obtain insights on the behavior of symmetric Dunkl processes and Dyson's model. Let us consider the effect of the intertwining operator on the quadratic equations $m_2(\boldsymbol{x}) = \sum_{i=1}^N x_i^2 = 1$ and

$m_{11}(\boldsymbol{x}) = \sum_{1\le i<j\le N} x_i x_j = 1$.

$$\begin{aligned} V_k \sum_{j=1}^N x_j^2 &= \frac{k+1}{kN+1}\sum_{j=1}^N x_j^2 + \frac{2k}{kN+1}\sum_{1\le i<j\le N} x_i x_j = 1 \\ V_k \sum_{1\le i<j\le N} x_i x_j &= \frac{k(N-1)}{2(kN+1)}\sum_{j=1}^N x_j^2 + \frac{k(N-1)+1}{kN+1}\sum_{1\le i<j\le N} x_i x_j = 1 \end{aligned} \tag{38}$$

From the coefficients of the sums above, it is clear that when $k=0$ the polynomials on the LHS are unchanged. When $k=1$, these equations become

$$\begin{aligned} V_{k=1} \sum_{j=1}^N x_j^2 &= \frac{2}{N+1}\left(\sum_{j=1}^N x_j^2 + \sum_{1\le i<j\le N} x_i x_j\right) = 1, \\ V_{k=1} \sum_{1\le i<j\le N} x_i x_j &= \frac{(N-1)}{2(N+1)}\sum_{j=1}^N x_j^2 + \frac{N}{N+1}\sum_{1\le i<j\le N} x_i x_j = 1. \end{aligned} \tag{39}$$

If we take the limit $k \to \infty$, the two quadratic equations become

$$\begin{aligned} V_k \sum_{j=1}^N x_j^2 &\xrightarrow{k\to\infty} \frac{1}{N}\sum_{j=1}^N x_j^2 + \frac{2}{N}\sum_{1\le i<j\le N} x_i x_j = \frac{1}{N}\left(\sum_{j=1}^N x_j\right)^2 = 1, \\ V_k \sum_{1\le i<j\le N} x_i x_j &\xrightarrow{k\to\infty} \frac{N-1}{2N}\sum_{j=1}^N x_j^2 + \frac{N-1}{N}\sum_{1\le i<j\le N} x_i x_j = \frac{N-1}{2N}\left(\sum_{j=1}^N x_j\right)^2 = 1. \end{aligned} \tag{40}$$

These equations represent two pairs of planes, namely

$$\sum_{j=1}^N x_j = \pm\sqrt{N} \quad \text{and} \quad \sum_{j=1}^N x_j = \pm\sqrt{\frac{2N}{N-1}}. \tag{41}$$

We plot Eqs. (38) for the particular case $N=3$ in Fig. 1. The line defined by the vector $(1,1,1)$ is represented by the diagonal line in each of the subfigures. From the figure, we can see that as the value of k increases, the surfaces get stretched equally in all directions orthogonal to the line. In the limit $k \to \infty$, the surfaces become planes, and there are two of them because we have chosen quadratic polynomials, i.e., $|\lambda| = 2$.

From these observations, we expect that the effect of $V_{k\to\infty}$ on the monomial symmetric functions m_λ is to reduce them to the $|\lambda|$-th power of $m_1(\boldsymbol{x}) = \sum_{j=1}^N x_j$. The power must be $|\lambda|$ because V_k conserves the degree of all homogeneous polynomials by definition. We prove this fact in the following theorem.



(a) $V_{k=0}\sum_{j=1}^{3}x_j^2=1$ (b) $V_{k=1}\sum_{j=1}^{3}x_j^2=1$ (c) $V_{k\to\infty}\sum_{j=1}^{3}x_j^2=1$



(d) $V_{k=0}\sum_{1\le i<j\le 3}x_ix_j=1$ (e) $V_{k=1}\sum_{1\le i<j\le 3}x_ix_j=1$ (f) $V_{k\to\infty}\sum_{1\le i<j\le 3}x_ix_j=1$

FIG. 1: Effect of the intertwining operator on the equations $m_2(\boldsymbol{x})=1$ and $m_{11}(\boldsymbol{x})=1$.

C. The $k\to\infty$ limit of Theorem 2

Theorem 3. *The limit $k\to\infty$ of Theorem 2 is*

$$\lim_{k\to\infty}V_k m_\lambda(\boldsymbol{x})=\frac{M(\lambda,N)}{N^{|\lambda|}}\left(\sum_{j=1}^{N}x_j\right)^{|\lambda|}. \tag{42}$$

Proof. The quantity

$$\frac{c_\tau(1/k)}{c'_\tau(1/k)(kN)_\tau^{(1/k)}}=\prod_{(i,j)\in\tau}\frac{(\tau_i-j+k(\tau'_j-i+1))}{(k(N-i+1)+j-1)(\tau_i-j+1+k(\tau'_j-i))}$$

tends to zero as $k\to\infty$ in general. The only case in which this quantity does not vanish is when $\tau'_j=i$ for any $(i,j)\in\tau$. Now, τ'_j represents the number of parts greater than or equal to j in τ (see Appendix A), so we can conclude that this condition is satisfied only when $\tau'_j=1$, i.e., $\tau=(\tau_1,0,\ldots)$. In other words, only partitions of length equal to one satisfy this condition.

Therefore,

$$\frac{c_\tau(1/k)}{c'_\tau(1/k)(kN)_\tau^{(1/k)}} \overset{k\to\infty}{\longrightarrow} 0$$

whenever $l(\tau) > 1$ and

$$\frac{c_\tau(1/k)}{c'_\tau(1/k)(kN)_\tau^{(1/k)}} = \prod_{j=1}^{\tau_1} \frac{(\tau_1 - j + k)}{(kN + j - 1)(\tau_1 - j + 1)} \overset{k\to\infty}{\longrightarrow} \prod_{j=1}^{\tau_1} \frac{1}{N(\tau_1 - j + 1)} = \frac{1}{N^{\tau_1}\tau_1!}$$

when $l(\tau) = 1$. Then, V_k becomes

$$V_k m_\lambda(\boldsymbol{x}) \overset{k\to\infty}{\longrightarrow} \frac{\lambda! M(\lambda, N)}{N^{|\lambda|}|\lambda|!} a_{|\lambda|,\lambda}(e_1(\boldsymbol{x}))^{|\lambda|},$$

due to the fact that if $l(\tau) = 1$, then $\tau' = (1, 1, \ldots, 1)$, a partition composed of ones with $|\lambda|$ parts (see Eqs. (A4) and (A6) for the definitions of $a_{|\lambda|,\lambda}$ and $e_1(\boldsymbol{x})$). We finish our calculation giving an explicit form of the matrix components $a_{|\lambda|,\lambda}$ by expanding $(e_1(\boldsymbol{x}))^{|\lambda|}$ in terms of $m_\tau(\boldsymbol{x})$. We write

$$(e_1(\boldsymbol{x}))^{|\lambda|} = \left(\sum_{j=1}^{N} x_j\right)^{|\lambda|} = \sum_{\substack{\tau: l(\tau)\leq N \\ |\tau| = |\lambda|}} \frac{|\lambda|!}{\tau!} m_\tau(\boldsymbol{x}) = \sum_{\substack{\tau: l(\tau)\leq N \\ |\tau| = |\lambda|}} a_{|\lambda|,\tau} m_\tau(\boldsymbol{x}),$$

and deduce that

$$a_{|\lambda|,\lambda} = \frac{|\lambda|!}{\lambda!}, \tag{43}$$

which proves the statement. □

Note that the equation

$$\frac{M(\lambda, N)}{N^{|\lambda|}} \left(\sum_{j=1}^{N} x_j\right)^{|\lambda|} = 1 \tag{44}$$

clearly represents one (for $|\lambda|$ odd) or two (for $|\lambda|$ even) planes with normal vector equal to

$$\mathbf{1} = (1, \ldots, 1). \tag{45}$$

This is consistent with Figs. 1(c) and (f).

V. ZERO-TEMPERATURE LIMIT OF DYSON'S MODEL

The physical implications of the observations in the previous section involve the behavior of the symmetric Dunkl process when k tends to infinity. This limit is equivalent to the zero-temperature limit ($\beta \to \infty$) of Dyson's model. Although the observations in the previous section suggest that the limit $\lim_{k\to\infty} \sum_{\rho\in S_N} E_k(\rho \boldsymbol{x}, \boldsymbol{y})$ exists, one must be careful when taking this limit in Eq. (7). In particular, the fact that $k = \beta/2$ is the coupling constant of the interaction between particles in Dyson's model implies that this interaction separates the particles infinitely fast when $k \to \infty$. In order to obtain an intuitive picture of what occurs at zero temperature, we consider the Kolmogorov forward equation associated to Eq. (4) with the parameter k in accordance with Theorem 1:

$$\begin{aligned} \frac{\partial}{\partial t} p_k^s(t, \boldsymbol{y}|\boldsymbol{x}) = \frac{1}{2}\Delta^{(y)} p_k^s(t, \boldsymbol{y}|\boldsymbol{x}) - \sum_{i=1}^{N} \sum_{\substack{j=1 \\ j\neq i}}^{N} \frac{k}{y_i - y_j} \frac{\partial}{\partial y_i} p_k^s(t, \boldsymbol{y}|\boldsymbol{x}) \\ + k \sum_{i=1}^{N} \sum_{\substack{j=1 \\ j\neq i}}^{N} \frac{p_k^s(t, \boldsymbol{y}|\boldsymbol{x})}{(y_i - y_j)^2}. \end{aligned} \tag{46}$$

We will now apply the scaling $\boldsymbol{y} \to \sqrt{k}\boldsymbol{v}$. Under this scaling, Eq. (46) becomes

$$\begin{aligned} \frac{\partial}{\partial t} p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x}) = \frac{1}{2k}\Delta^{(v)} p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x}) - \sum_{i=1}^{N} \sum_{\substack{j=1 \\ j\neq i}}^{N} \frac{1}{v_i - v_j} \frac{\partial}{\partial v_i} p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x}) \\ + \sum_{i=1}^{N} \sum_{\substack{j=1 \\ j\neq i}}^{N} \frac{p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x})}{(v_i - v_j)^2}. \end{aligned} \tag{47}$$

Furthermore, if we assume that $\lim_{k\to\infty} p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x}) k^{N/2} = p_\infty^s(t, \boldsymbol{v}|\boldsymbol{x})$, taking the limit $k \to \infty$ transforms Eq. (47) into

$$\frac{\partial}{\partial t} p_\infty^s(t, \boldsymbol{v}|\boldsymbol{x}) = -\sum_{i=1}^{N} \frac{\partial}{\partial v_i} \left[\sum_{\substack{j=1 \\ j\neq i}}^{N} \frac{p_\infty^s(t, \boldsymbol{v}|\boldsymbol{x})}{v_i - v_j} \right], \tag{48}$$

a Kolmogorov forward equation with a drift term but without a diffusion term. This equation suggests that the scaled process on $\boldsymbol{v}$ is actually deterministic. Physically speaking, this is a natural consequence of taking the $k \to \infty$ limit, since the randomness of Dyson's model is removed at the zero-temperature limit. Therefore, the argument above implies that $p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x}) k^{N/2}$ should converge to a function of the form

$$\lim_{k\to\infty} p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x}) k^{N/2} = \delta^N[\boldsymbol{v} - \boldsymbol{F}(t, \boldsymbol{x})], \tag{49}$$

where $\boldsymbol{F}(t, \boldsymbol{x})$ denotes the vector solution of a deterministic equation of motion.

Interestingly, the evolution of the scaled process considered above is related to the Hermite polynomials defined by [46]

$$H_N(x) = (-1)^N e^{x^2} \frac{\mathrm{d}^N}{\mathrm{d}x^N}(e^{-x^2}). \tag{50}$$

More specifically, the process described by Eq. (48) depends on the N roots of $H_N(x)$, here denoted by

$$\boldsymbol{z}_N = (z_{1,N}, \ldots, z_{N,N}), \tag{51}$$

and their permutations $\rho \boldsymbol{z}_N$, $\rho \in S_N$, where $z_{i,N}$ is the i-th root and $z_{i,N} < z_{j,N}$ for $i < j$. These N roots are known to be real [47]. Also, because the Hermite polynomials obey the relation

$$H_N(-x) = (-1)^N H_N(x), \tag{52}$$

their roots add up to zero, i.e.,

$$\sum_{j=1}^{N} z_{j,N} = 0. \tag{53}$$

We address the zero-temperature limit in the following theorem.

Theorem 4. *The $k \to \infty$ limit of the TPD of the symmetric Dunkl process $p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x})k^{N/2}$ is independent of $\boldsymbol{x}$ and is given by*

$$p_\infty^s(t, \boldsymbol{v}|\boldsymbol{x}) = p_\infty^s(t, \boldsymbol{v}) = \sum_{\rho \in S_N} \delta^N[\boldsymbol{v} - \sqrt{2t}\rho \boldsymbol{z}_N]. \tag{54}$$

Proof. We first calculate the limit

$$\lim_{k\to\infty} \sum_{\rho \in S_N} E_k(\rho \boldsymbol{x}, \boldsymbol{y}). \tag{55}$$

Using Theorems 2 and 3 and the expansion in Eq. (33), we obtain

$$\begin{aligned}
\lim_{k\to\infty} \sum_{\rho \in S_N} E_k(\rho \boldsymbol{x}, \boldsymbol{y}) &= \sum_{\mu : l(\mu) \le N} \frac{N! m_\mu(\boldsymbol{y})}{\mu! M(\mu, N)} \lim_{k\to\infty} V_k m_\mu(\boldsymbol{x}) \\
&= \sum_{\mu : l(\mu) \le N} \frac{N! m_\mu(\boldsymbol{y})}{\mu! N^{|\mu|}} \left(\sum_{j=1}^{N} x_j \right)^{|\mu|} \\
&= \sum_{\mu : l(\mu) \le N} \frac{N!}{\mu!} m_\mu \left(\frac{(\boldsymbol{x} \cdot \mathbf{1})\boldsymbol{y}}{N} \right) \\
&= N! \exp\left(\frac{(\boldsymbol{x} \cdot \mathbf{1})(\boldsymbol{y} \cdot \mathbf{1})}{N} \right).
\end{aligned} \tag{56}$$

We now turn our attention to $p_k^s(t, \boldsymbol{y}|\boldsymbol{x})$. The probability that the symmetric Dunkl process reaches an infinitesimal volume $\mathrm{d}^N y$ around $\boldsymbol{y}$ from $\boldsymbol{x}$ after a time t is $p_k^s(t, \boldsymbol{y}|\boldsymbol{x})\, \mathrm{d}^N y$. This probability, after the scaling $\boldsymbol{y} \to \sqrt{k}\boldsymbol{v}$ equals

$$p_k^s(t, \boldsymbol{y}|\boldsymbol{x})\, \mathrm{d}^N y = p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x}) k^{N/2}\, \mathrm{d}^N v = \exp\left\{ \log\left(p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x}) k^{N/2} \right) \right\} \mathrm{d}^N v. \tag{57}$$

When k is sufficiently large one can use Stirling's approximation [48] combined with Eqs. (13), (28) and (56) to approximate the probability density in Eq. (57) as the exponential of

$$\begin{aligned} &k\left[\frac{N}{2}(N-1)(1-\log t)-\sum_{j=1}^{N} j\log j+2\log|h_N(\boldsymbol{v})|-\frac{v^2}{2t}\right] \\ &+\frac{\sqrt{k}}{N}(\boldsymbol{x}\cdot\mathbf{1})(\boldsymbol{v}\cdot\mathbf{1})+\frac{N}{2}\log k+\log(N!)-\frac{N}{2}\log(2\pi t)-\frac{x^2}{2t}. \end{aligned} \tag{58}$$

Let us denote the term in the square brackets by $F_N(\boldsymbol{v},t)$. According to this expression, in the limit $k\to\infty$ the probability density goes to infinity whenever $F_N(\boldsymbol{v},t)$ is positive, and vanishes when it is negative. The lower-order terms in k are negligible in either of these two cases. However, this distinction is difficult to make due to the presence of the Vandermonde determinant of $\boldsymbol{v}$ in $F_N(\boldsymbol{v},t)$. We show in Appendix F that $F_N(\boldsymbol{v},t)$ is negative except in the points

$$\boldsymbol{v}=\sqrt{2t}\rho\boldsymbol{z}_N. \tag{59}$$

The consequence of this is that the probability density (57) vanishes at $\boldsymbol{v}\neq\sqrt{2t}\boldsymbol{z}_N$ or its permutations. On the other hand, when $\boldsymbol{v}=\sqrt{2t}\boldsymbol{z}_N$ or its permutations, $kF_N(\boldsymbol{v},t)$ (see Appendix F) and the term proportional to $\sqrt{k}$ vanish, because the roots of the Hermite polynomials add up to zero. That is,

$$\frac{\sqrt{k}}{N}(\boldsymbol{x}\cdot\mathbf{1})(\sqrt{2t}\boldsymbol{z}_N\cdot\mathbf{1})=\frac{\sqrt{2kt}}{N}(\boldsymbol{x}\cdot\mathbf{1})\sum_{i=1}^{N} z_{i,N}=0. \tag{60}$$

The only k-dependent term that remains is $\frac{N}{2}\log(k)$, which tends to infinity with k. Therefore, we can write

$$\lim_{k\to\infty} p_k^s(t,\sqrt{k}\boldsymbol{v}|\boldsymbol{x})k^{N/2}\,\mathrm{d}^N v\propto\sum_{\rho\in S_N}\delta[\boldsymbol{v}(t)-\sqrt{2t}\rho\boldsymbol{z}_N]. \tag{61}$$

However, we know that $p_k^s(t,\sqrt{k}\boldsymbol{v}|\boldsymbol{x})k^{N/2}\,\mathrm{d}^N v$ is equal to $N!$ when integrated over $\mathbb{R}^N$, for any value of k. Similarly, the integral over $\mathbb{R}^N$ of the sum of delta functions on the RHS is also equal to $N!$. Therefore, the two quantities above are equal. □

Note that the effect of $\lim_{k\to\infty}\sum_{\rho\in S_N}E_k(\rho\boldsymbol{x},\boldsymbol{y})$ on the final result is that it makes the TPD $p_\infty^s(t,\boldsymbol{v}|\boldsymbol{x})$ independent of $\boldsymbol{x}$. When $F_N(\boldsymbol{v},t)$ attains its maximum value, the symmetric Dunkl kernel is equal to one for all values of $\boldsymbol{x}$, which leaves $k^{N/2}$ as the dominating term in the limit. Therefore, the form of the intertwining operator and the properties of the Hermite polynomials are ultimately responsible for removing the dependence of p_k^s on $\boldsymbol{x}$ in the zero-temperature limit.

VI. CONCLUDING REMARKS

We established the equivalence between Dyson's model and symmetric Dunkl processes under the parameter relation (14) in Theorem 1. This correspondence yields a physical interpretation of the abstract parameter k. The relation $\beta/2 = k$ implies that one can give to k the interpretation of inverse temperature given to the parameter β. We have also extracted the effect of the intertwining operator on symmetric polynomials in Theorem 2 using Theorem 1. It is of interest to note that the functions involved, which are commonly found in combinatorics and representation theory [28, 29] and in quantum integrable systems [32–34], are found here through the comparison of two multivariate stochastic processes. This result stresses the importance of the role that symmetric polynomials play in the setting of stochastic processes. While it is known that Schur functions are essential in the study of the non-colliding Brownian motion ($\beta/2 = k = 1$) [7, 9, 26, 27], their role in the general k case is played by Jack functions [4, 43], which makes them equally essential for the study of Dyson's model. We showed in Theorem 3 that the limit $k \to \infty$ of Theorem 2 exists, and that in this limit V_k has a simplified form. This fact allowed us to calculate the zero-temperature limit of the symmetrized Dunkl kernel, and in turn obtain the TPD of the symmetric Dunkl process, which is the result of Theorem 4. The positions of the Brownian particles are locked to a deterministic path, which is a natural consequence of the lack of thermal vibrations in this regime.

Moreover, we pointed out the following two properties of the symmetric Dunkl process in the limit $k \to \infty$ in Theorem 4. The first of them is that the final configuration of the Brownian particles depends on the roots of the Hermite polynomials rather than being an equally-spaced or simpler configuration. This is a consequence of the form of the interaction between particles: the Vandermonde determinant of $\boldsymbol{v}$ appears in $p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x})$ as a death factor that makes the TPD vanish whenever $v_i = v_j,\ i \neq j$, pushing the particles away from each other. Furthermore, the partial derivatives of the Vandermonde determinant are responsible for the form of the drift term in Eqs. (4) and (10). Therefore, a different interaction would lead to a different form for the function $F_N(\boldsymbol{v}, t)$, which would in turn have its maxima in points different from $\boldsymbol{z}_N$ or its permutations.

The second is that in the zero-temperature limit, $p_k^s(t, \sqrt{k}\boldsymbol{v}|\boldsymbol{x})k^{N/2}$ is independent of the initial configuration $\boldsymbol{x}$. Because the process is deterministic in this limit, one expects it to depend on the initial conditions from which it has evolved. Contrary to this expectation, the nature of the intertwining operator at $k \to \infty$ and the symmetry that the roots of the Hermite polynomials obey take away that dependence.

In summary, we have succeeded in extracting an expression for the intertwining operator, we have observed its effect on symmetric polynomials and we have used it to extract information about Dyson's model and its behavior at zero temperature. As mentioned in Secs. I and II A, random matrix theory and its stochastic version, Dyson's model, have been thoroughly studied and applied in numerous branches of physics and mathematics. Therefore, in view of the present results we expect Dunkl processes and Dunkl theory to be applied more broadly in physics in general, and in non-equilibrium statistical mechanics in particular.

There are several aspects of Dunkl processes that require further study. An expression for the intertwining operator for non-symmetric polynomials is desired in order to obtain the TPD of non-symmetric A-type Dunkl processes. This, in turn, would open the possibility of studying closely the nature of the spontaneous jumps that these processes undergo due to the jump term in their Kolmogorov backward equation (the third term on the RHS of Eq. (10)). Finally, the existence of explicit forms for the symmetrized Dunkl kernel (the LHS of Eq. (29)) for the B and D-type root systems as given in Refs. [43, 49] should allow for other types of Dunkl processes to be applied in physical systems where symmetry with respect to those root systems is required.

ACKNOWLEDGMENTS

SA would like to thank Takashi Mori for his helpful insights on this work. SA is supported by the MEXT scholarship for research students. MK is supported by the Grant-in-Aid for Scientific Research (C) (Grant No.21540397) from Japan Society for the Promotion of Science.

Appendix A: Multivariate Special Functions

We begin by listing briefly the definitions involving integer partitions that are necessary to properly define the multivariate functions used in Secs. IV and V [29]. A partition τ of an integer n is a non-negative integer vector such that its components are arranged in decreasing order while adding up to n. The components of τ are called parts, and its length, $l(\tau)$, is the number of non-zero parts in it. The sum of the parts is called the modulus of the partition and is denoted by $|\tau|$. In other words, τ is an integer partition of n if $\tau_i \geq \tau_j$ for $i < j$, and

$$|\tau| = \sum_{i=1}^{l(\tau)} \tau_i = n. \tag{A1}$$

For example, $\tau = (\tau_1, \tau_2, \ldots) = (4, 4, 2, 0, \ldots)$ is an integer partition of $n = 10$ and has length 3. Additionally, the symbol $\tau!$ denotes the product $\prod_{j=1}^{l(\tau)} \tau_j!$. We will also use the natural ordering for partitions of the same integer defined by

$$\lambda < \tau \;\Leftrightarrow\; \lambda_1 + \ldots + \lambda_j \leq \tau_1 + \ldots + \tau_j \;{}^{\forall} j = 1, \ldots, N, \tag{A2}$$

where λ and τ are assumed to be different partitions.

We say that the ordered pair (i, j) is in the partition τ (i.e., $(i, j) \in \tau$) when $1 \leq i \leq l(\tau)$ and $1 \leq j \leq \tau_i$. The number τ_j' is the number of parts τ_l such that $\tau_l \geq j$. Finally, the partition formed by the τ_j' is denoted by τ', and it is called the conjugate partition of τ.

Let us now introduce three families of symmetric polynomials, keeping in mind that we will consider polynomials of N variables. Given an integer partition τ, the monomial symmetric functions are given by

$$m_\tau(\boldsymbol{x}) = \sum_{\sigma} \prod_{j=1}^{N} x_j^{\tau_{\sigma(j)}}, \tag{A3}$$

where the sum is taken over all permutations σ such that each monomial $\prod_{j=1}^{N} x_j^{\tau_{\sigma(j)}}$ is distinct.

For an integer $0 \leq n \leq N$, the elementary symmetric function $e_n(\boldsymbol{x})$ is given by

$$e_n(\boldsymbol{x}) = \sum_{1 \leq i_1 < \cdots < i_n \leq N} \prod_{j=1}^{n} x_{i_j}. \tag{A4}$$

For example, $e_0(\boldsymbol{x}) = 1$, $e_1(\boldsymbol{x}) = \sum_{i=1}^{N} x_i$, $e_2(\boldsymbol{x}) = \sum_{1 \leq i < j \leq N} x_i x_j$ and $e_N(\boldsymbol{x}) = \prod_{i=1}^{N} x_i$. When the subscript of e is a partition, it is given by

$$e_\tau(\boldsymbol{x}) = \prod_{i=1}^{l(\tau)} e_{\tau_i}(\boldsymbol{x}). \tag{A5}$$

It is known that the relationship between the monomial and elementary symmetric functions is given by

$$e_{\tau'}(\boldsymbol{x}) = \sum_{\substack{\lambda : \lambda \leq \tau \\ |\lambda| = |\tau|}} a_{\tau\lambda} m_\lambda(\boldsymbol{x}), \tag{A6}$$

where $a_{\tau\lambda}$ is an upper triangular matrix with diagonal elements equal to one. Note that the subscript on the LHS is the conjugate partition of τ.

The Schur function $s_\tau(\boldsymbol{x})$ is given by the relation

$$s_\tau(\boldsymbol{x}) = m_\tau(\boldsymbol{x}) + \sum_{\substack{\lambda : \lambda < \tau \\ |\lambda| = |\tau|}} K_{\tau\lambda} m_\lambda(\boldsymbol{x}) = \frac{\det_{1 \leq i,j \leq N}[x_j^{\tau_i + N - i}]}{\det_{1 \leq i,j \leq N}[x_j^{N-i}]}. \tag{A7}$$

(The second equality is the Jacobi-Trudi formula [28, 29]). The matrix $K_{\tau\lambda}$ is called the Kostka matrix.

The Jack function of parameter $\alpha > 0$, $\mathcal{C}_\tau^{(\alpha)}(\boldsymbol{x})$ is defined as the polynomial eigenfunction of the operator [50]

$$\left(\sum_{i=1}^N x_i^2 \frac{\partial^2}{\partial x_i^2} + \frac{2}{\alpha} \sum_{1\le i\ne j\le N} \frac{x_i^2}{x_i - x_j}\frac{\partial}{\partial x_i}\right) \mathcal{C}_\tau^{(\alpha)}(\boldsymbol{x}) = E_{\tau,\alpha}\mathcal{C}_\tau^{(\alpha)}(\boldsymbol{x}) \tag{A8}$$

with eigenvalue

$$E_{\tau,\alpha} = \sum_{j=1}^N \tau_j(\tau_j - 1 - 2(j-1)/\alpha) + |\tau|(N-1). \tag{A9}$$

There is one Jack function for each partition τ, which is unique up to a normalization constant. We will use the following two normalizations: the C normalization, which is defined so that

$$\sum_{n=0}^\infty \frac{1}{n!} \sum_{\substack{\tau:l(\tau)\le N\\|\tau|=n}} \mathcal{C}_\tau^{(\alpha)}(\boldsymbol{x}) = \exp(x_1 + x_2 + \ldots + x_N) \tag{A10}$$

for all α, and the P normalization, which is defined by the following linear combination of the monomial symmetric functions $m_\lambda(\boldsymbol{x})$,

$$\mathcal{P}_\tau^{(\alpha)}(\boldsymbol{x}) = m_\tau(\boldsymbol{x}) + \sum_{\substack{\lambda:\lambda<\tau\\|\lambda|=|\tau|}} u_{\tau\lambda}(\alpha) m_\lambda(\boldsymbol{x}). \tag{A11}$$

The partition-indexed matrix $u_{\tau\lambda}(\alpha)$, as shown in Ref. [29], is an upper triangular matrix whose diagonal entries are equal to one, and its non-diagonal entries are sums of ratios of the form $(a\alpha + b)/(c\alpha + d)$, where a, b, c and d are non-negative integers. Therefore, it can be inverted, yielding the inverse relation

$$m_\tau(\boldsymbol{y}) = \sum_{\substack{\lambda:\lambda\le\tau\\|\lambda|=|\tau|}} (u^{-1})_{\tau\lambda}(\alpha)\mathcal{P}_\lambda^{(\alpha)}(\boldsymbol{y}). \tag{A12}$$

Also, the limits $\lim_{\alpha\to 0} u_{\tau\lambda}(\alpha)$ and $\lim_{\alpha\to\infty} u_{\tau\lambda}(\alpha)$ are well-defined. In fact, it is known that [29]

$$\begin{aligned} \mathcal{P}_\tau^{(0)}(\boldsymbol{x}) &= e_{\tau'}(\boldsymbol{x}) \\ \mathcal{P}_\tau^{(1)}(\boldsymbol{x}) &= s_\tau(\boldsymbol{x}) \\ \mathcal{P}_\tau^{(\infty)}(\boldsymbol{x}) &= m_\tau(\boldsymbol{x}), \end{aligned} \tag{A13}$$

or, equivalently,

$$\begin{aligned} \lim_{\alpha\to 0} u_{\tau\lambda}(\alpha) &= a_{\tau\lambda} \\ u_{\tau\lambda}(1) &= K_{\tau\lambda} \\ \lim_{\alpha\to\infty} u_{\tau\lambda}(\alpha) &= \delta_{\tau\lambda}. \end{aligned} \tag{A14}$$

Using Jack functions, one can define the generalized hypergeometric function ${}_0\mathcal{F}_0^{(\alpha)}(\boldsymbol{x}, \boldsymbol{y})$ as follows [4, 43],

$$ {}_0\mathcal{F}_0^{(\alpha)}(\boldsymbol{x}, \boldsymbol{y}) = \sum_{n=0}^{N} \frac{1}{n!} \sum_{\substack{\tau : l(\tau) \leq N \\ |\tau| = n}} \frac{\mathcal{C}_\tau^{(\alpha)}(\boldsymbol{x}) \mathcal{C}_\tau^{(\alpha)}(\boldsymbol{y})}{\mathcal{C}_\tau^{(\alpha)}(\boldsymbol{1})}, \tag{A15} $$

with $\boldsymbol{1}$ as given in Eq. (45). For the general definition of the generalized hypergeometric function ${}_p\mathcal{F}_q^{(\alpha)}$, see Ref. [43]. It is desirable to express the above in terms of the P-normalized Jack functions so that we can apply Eqs. (A13) and (A14). For this purpose, we require the generalized Pochhammer symbol $(a)_\tau^{(\alpha)}$, which is defined as the product

$$ (a)_\tau^{(\alpha)} = \prod_{i=1}^{l(\tau)} \frac{\Gamma(a - (i-1)/\alpha + \tau_i)}{\Gamma(a - (i-1)/\alpha)}, \tag{A16} $$

as well as the functions

$$ \begin{aligned} c_\tau(\alpha) &= \prod_{(i,j)\in\tau} (\alpha(\tau_i - j) + \tau'_j - i + 1), \\ c'_\tau(\alpha) &= \prod_{(i,j)\in\tau} (\alpha(\tau_i - j + 1) + \tau'_j - i). \end{aligned} \tag{A17} $$

With these definitions, one can write [51]

$$ \begin{aligned} \mathcal{C}_\tau^{(\alpha)}(\boldsymbol{x}) &= \frac{\alpha^{|\tau|}|\tau|!}{c'_\tau(\alpha)} \mathcal{P}_\tau^{(\alpha)}(\boldsymbol{x}), \\ \mathcal{P}_\tau^{(\alpha)}(\boldsymbol{1}) &= \frac{\alpha^{|\tau|}(N/\alpha)_\tau^{(\alpha)}}{c_\tau(\alpha)}. \end{aligned} \tag{A18} $$

Insertion of the above in Eq. (A15) yields, with $\alpha = 1/k$,

$$ {}_0\mathcal{F}_0^{(1/k)}(\boldsymbol{x}, \boldsymbol{y}) = \sum_{n=0}^{N} \sum_{\substack{\tau : l(\tau) \leq N \\ |\tau| = n}} \frac{c_\tau(1/k)}{c'_\tau(1/k)} \frac{\mathcal{P}_\tau^{(1/k)}(\boldsymbol{x}) \mathcal{P}_\tau^{(1/k)}(\boldsymbol{y})}{(kN)_\tau^{(1/k)}}. \tag{A19} $$

Appendix B: Root Systems

The definition of a root system depends on the reflection operator [37], which is given by

$$ \sigma_{\boldsymbol{\alpha}} \boldsymbol{x} = \boldsymbol{x} - 2 \frac{\boldsymbol{x} \cdot \boldsymbol{\alpha}}{\boldsymbol{\alpha} \cdot \boldsymbol{\alpha}} \boldsymbol{\alpha}, \tag{B1} $$

where $\boldsymbol{\alpha}, \boldsymbol{x} \in \mathbb{R}^N$, and $\sigma_{\boldsymbol{\alpha}} \boldsymbol{x}$ denotes the vector obtained by reflecting $\boldsymbol{x}$ across the plane normal to the vector $\boldsymbol{\alpha}$.

A root system is defined as a finite set of vectors R which is invariant under reflections along its own elements, called roots. That is, $\sigma_{\boldsymbol{\alpha}}\boldsymbol{\xi} \in R$ for all $\boldsymbol{\alpha}, \boldsymbol{\xi} \in R$. R has the property that, for any one of its elements $\boldsymbol{\xi}$, $b\boldsymbol{\xi} \in R$ implies $b = \pm 1$. It is assumed that none of the roots is a zero vector.

For every root system, a base can be chosen such that all roots are a linear combination of the base vectors with either all coefficients negative or all coefficients positive. The elements of such a base are called simple roots, and they divide the root system into the positive and negative subsystems, here denoted as R_+ and R_-. The choice of simple roots is not unique, but the positive and negative subsystems they define always divide the root system into two disjoint parts of the same size (cardinality).

The multiplicity function k is a function of a root in R that returns a real value and that is invariant under reflections along elements of R. That is,

$$k(\boldsymbol{\alpha}) = k(\sigma_{\boldsymbol{\alpha}'}\boldsymbol{\alpha}) \ {}^{\forall}\boldsymbol{\alpha}, \boldsymbol{\alpha}' \in R. \tag{B2}$$

The multiplicity function can be understood as a set of parameters (or multiplicities) which are assigned to each subset of R formed by roots that are related by a reflection (or composition of reflections) along some other root(s).

It is a well-known fact [37] that the multiplicity function reduces to a single parameter for the A_{N-1} root system. For simplicity, we will denote this root system by A. Let us prove this fact as follows. A is given by

$$A = \{\boldsymbol{\alpha}_{ij} = (\boldsymbol{e}_i - \boldsymbol{e}_j)/\sqrt{2} : i, j = 1, \dots, N; i \neq j\}. \tag{B3}$$

We will choose the positive subsystem

$$A_+ = \{\boldsymbol{\alpha}_{ij} = (\boldsymbol{e}_i - \boldsymbol{e}_j)/\sqrt{2} : 1 \leq j < i \leq N\}, \tag{B4}$$

where $\boldsymbol{e}_i$ denotes the i-th unit base vector. This positive subsystem is generated by the simple roots

$$\boldsymbol{\alpha}_{i+1,i} = (\boldsymbol{e}_{i+1} - \boldsymbol{e}_i)/\sqrt{2}, \ i = 1, \dots, N-1. \tag{B5}$$

Hereafter, we will write $\sigma_{\alpha_{ij}} = \sigma_{ij}$. We must note that the effect of σ_{ij} on an arbitrary vector $\boldsymbol{x}$ is that of exchanging its i-th and j-th components. To see this, we will compute the l-th component of $\sigma_{ij}\boldsymbol{x}$:

$$(\sigma_{ij}\boldsymbol{x})_l = x_l - (x_i - x_j)(\delta_{il} - \delta_{jl}). \tag{B6}$$

It is easy to see that x_l remains unchanged for $l \neq i, j$, and that $(\sigma_{ij}\boldsymbol{x})_i = x_j$ and $(\sigma_{ij}\boldsymbol{x})_j = x_i$. Therefore, the group generated by the reflections along the elements of A with composition as the group operation is the symmetric group S_N.

In view of this property of A, we see that we can obtain any root from at most two reflections of any other root, as shown below. Consider an arbitrary root $\boldsymbol{\alpha}_{ij}$ and apply to it the reflection σ_{mj}, with m arbitrary. This reflection exchanges the j-th and the m-th components of $\boldsymbol{\alpha}_{ij}$, leaving us with $\boldsymbol{\alpha}_{im}$. If we reflect once more using σ_{il}, with l arbitrary, we obtain $\boldsymbol{\alpha}_{lm}$ as desired. Since k is invariant under any of these reflections, one obtains

$$k(\boldsymbol{\alpha}_{ij}) = k(\sigma_{mj}\sigma_{il}\boldsymbol{\alpha}_{ij}) = k(\boldsymbol{\alpha}_{lm})$$

in general, and therefore it can be concluded that k is independent of its argument, so it is a single parameter.

Appendix C: Dunkl Operators and Dunkl Heat Equation

Dunkl operators are given by [30],

$$T_{\boldsymbol{\xi}} f(\boldsymbol{x}) = \partial_{\boldsymbol{\xi}} f(\boldsymbol{x}) + \sum_{\boldsymbol{\alpha}\in R_+} k(\boldsymbol{\alpha}) \frac{f(\boldsymbol{x}) - f(\sigma_{\boldsymbol{\alpha}}\boldsymbol{x})}{\boldsymbol{\alpha}\cdot\boldsymbol{x}} \boldsymbol{\alpha}\cdot\boldsymbol{\xi}. \tag{C1}$$

The first term is a derivative along the vector $\boldsymbol{\xi}$. Each of the summands in the second term is proportional to the odd part of f along the vector $\boldsymbol{\alpha}$.

Dunkl defined this operator in order to study multivariate orthogonal polynomials and special functions related to reflection groups. Once a root system and multiplicity function are chosen, the operator $T_{\boldsymbol{\xi}}$ commutes with $T_{\boldsymbol{\xi}'}$ and therefore has many of the characteristics of partial derivatives. Hereafter we will use the notation $T_i = T_{\boldsymbol{e}_i}$ as in Section II.

In Ref. [36], Rösler found the TPD $p_k(t, \boldsymbol{y}|\boldsymbol{x})$ that solves Eq. (9) as its Kolmogorov backward equation, with T_i given by Eq. (C1). The generalized heat equation (9), or Dunkl heat equation, is given explicitly by [30]

$$\begin{aligned}\frac{\partial}{\partial t} p_k(t, \boldsymbol{y}|\boldsymbol{x}) = \frac{1}{2}\Delta^{(x)} p_k(t, \boldsymbol{y}|\boldsymbol{x}) + \sum_{\boldsymbol{\alpha}\in R_+} k(\boldsymbol{\alpha}) \frac{\partial_{\boldsymbol{\alpha}} p_k(t, \boldsymbol{y}|\boldsymbol{x})}{\boldsymbol{\alpha}\cdot\boldsymbol{x}} \\ - \sum_{\boldsymbol{\alpha}\in R_+} k(\boldsymbol{\alpha}) \frac{\alpha^2}{2} \frac{p_k(t, \boldsymbol{y}|\boldsymbol{x}) - p_k(t, \boldsymbol{y}|\sigma_{\boldsymbol{\alpha}}\boldsymbol{x})}{(\boldsymbol{\alpha}\cdot\boldsymbol{x})^2}.\end{aligned} \tag{C2}$$

Appendix D: Dunkl Transform and Transition Probability Density

The definition of the Dunkl transform requires both the Dunkl kernel defined in Section III and the weight function

$$w_k(\boldsymbol{x}) = \prod_{\boldsymbol{\alpha}\in R} |\boldsymbol{\alpha}\cdot\boldsymbol{x}|^{k(\boldsymbol{\alpha})}. \tag{D1}$$

For fixed R and k and functions $f \in L^1(\mathbb{R}^N, w_k)$, the set of integrable functions with respect to the weight function $w_k(\boldsymbol{x})$, the Dunkl transform is defined by the equation [35]

$$\hat{f}_k(\boldsymbol{\xi}) = \frac{1}{c_k}\int_{\mathbb{R}^N} f(\boldsymbol{x}) E_k(-i\boldsymbol{\xi}, \boldsymbol{x}) w_k(\boldsymbol{x})\,\mathrm{d}^N x, \tag{D2}$$

and

$$c_k = \int_{\mathbb{R}^N} e^{-\xi^2/2} w_k(\boldsymbol{\xi})\,\mathrm{d}^N \xi \tag{D3}$$

is a normalization constant obtained from a Selberg integral [3]. Due to the properties of the Dunkl kernel it is easy to see that

$$T_j^{\boldsymbol{\xi}} \hat{f}_k(\boldsymbol{\xi}) = \frac{1}{c_k}\int_{\mathbb{R}^N} [-ix_j f(\boldsymbol{x})] E_k(-i\boldsymbol{\xi}, \boldsymbol{x}) w_k(\boldsymbol{x})\,\mathrm{d}^N x. \tag{D4}$$

On the other hand, since

$$\frac{\partial}{\partial x_j} w_k(\boldsymbol{x}) = 2\sum_{\boldsymbol{\alpha}\in R_+} \alpha_j \frac{k(\boldsymbol{\alpha})}{\boldsymbol{\alpha}\cdot\boldsymbol{x}} w_k(\boldsymbol{x}), \tag{D5}$$

a direct calculation shows that

$$\int_{\mathbb{R}^N} [T_j f(\boldsymbol{x})] g(\boldsymbol{x}) w_k(\boldsymbol{x})\,\mathrm{d}^N x = -\int_{\mathbb{R}^N} f(\boldsymbol{x}) [T_j g(\boldsymbol{x})] w_k(\boldsymbol{x})\,\mathrm{d}^N x \tag{D6}$$

for f and g integrable with respect to w_k. Using Eq. (D6), one can prove directly that

$$\frac{1}{c_k}\int_{\mathbb{R}^N} [T_j^{\boldsymbol{x}} f(\boldsymbol{x})] E_k(-i\boldsymbol{\xi}, \boldsymbol{x}) w_k(\boldsymbol{x})\,\mathrm{d}^N x = i\xi_j \hat{f}_k(\boldsymbol{\xi}). \tag{D7}$$

The inverse Dunkl transform is defined as

$$f(\boldsymbol{x}) = \frac{1}{c_k}\int_{\mathbb{R}^N} \hat{f}_k(\boldsymbol{\xi}) E_k(i\boldsymbol{\xi}, \boldsymbol{x}) w_k(\boldsymbol{\xi})\,\mathrm{d}^N \xi, \tag{D8}$$

and it is known that the inverse transform of the transform of f is equal to f almost everywhere. In view of the identity

$$\frac{1}{(2\pi)^{N/2}}\int_{\mathbb{R}^N} \hat{f}(\boldsymbol{\xi}) e^{i\boldsymbol{x}\cdot\boldsymbol{\xi}} e^{i\boldsymbol{y}\cdot\boldsymbol{\xi}}\,\mathrm{d}^N \xi = f(\boldsymbol{x}+\boldsymbol{y}), \tag{D9}$$

where $\hat{f}$ is the N-dimensional Fourier transform of f, the generalized Dunkl translation $\tau_{\boldsymbol{y}}$ is defined as

$$\tau_{\boldsymbol{y}} f(\boldsymbol{x}) = \frac{1}{c_k} \int_{\mathbb{R}^N} \hat{f}_k(\boldsymbol{\xi}) E_k(i\boldsymbol{x}, \boldsymbol{\xi}) E_k(i\boldsymbol{y}, \boldsymbol{\xi}) w_k(\boldsymbol{\xi}) \, \mathrm{d}^N \xi, \tag{D10}$$

which reduces to Eq. (D9) when $k = 0$. Because Eq. (D10) does not change when $\boldsymbol{x}$ and $\boldsymbol{y}$ are exchanged, we see that $\tau_{\boldsymbol{y}} f(\boldsymbol{x}) = \tau_{\boldsymbol{x}} f(\boldsymbol{y})$, and because the form of Eq. (D10) resembles that of Eq. (D8), we can write

$$\frac{1}{c_k} \int_{\mathbb{R}^N} \tau_{\boldsymbol{y}} f(\boldsymbol{x}) E_k(-i\boldsymbol{\xi}, \boldsymbol{x}) w_k(\boldsymbol{x}) \, \mathrm{d}^N x = \hat{f}_k(\boldsymbol{\xi}) E_k(i\boldsymbol{y}, \boldsymbol{\xi}). \tag{D11}$$

Finally, we list a useful consequence of Theorem 3.10 in Ref. [42] which is valid for $\boldsymbol{y}, \boldsymbol{z} \in \mathbb{C}^N$:

$$\frac{1}{c_k} \int_{\mathbb{R}^N} E_k(\boldsymbol{x}, \boldsymbol{y}) E_k(\boldsymbol{x}, \boldsymbol{z}) e^{-x^2/2} w_k(\boldsymbol{x}) \, \mathrm{d}^N x = e^{(y^2+z^2)/2} E_k(\boldsymbol{y}, \boldsymbol{z}). \tag{D12}$$

Now we will use the Dunkl kernel, transform and their properties to find the Green function Γ_k of the Dunkl heat equation (C2). First, we find a non-trivial solution to Eq. (C2) in $L^1(\mathbb{R}^N, w_k)$, which we denote $g_k(\boldsymbol{x}, t)$. Applying the Dunkl transform on Eq. (C2) and using Eq. (D7), we have

$$\frac{\partial}{\partial t} \hat{g}_k(\boldsymbol{\xi}, t) = -\frac{1}{2} \sum_{i=1}^{N} \xi_i^2 \hat{g}_k(\boldsymbol{x}, t) = -\frac{\xi^2}{2} \hat{g}_k(\boldsymbol{x}, t), \tag{D13}$$

where $\hat{g}_k(\boldsymbol{\xi}, t)$ is the Dunkl transform of $g_k(\boldsymbol{x}, t)$, and integrating with respect to t we obtain

$$\hat{g}_k(\boldsymbol{\xi}, t) = A e^{-t\xi^2/2}, \tag{D14}$$

where A is a constant independent of t. Let us define the sum of $k(\boldsymbol{\alpha})$ over the positive subsystem as

$$\gamma = \sum_{\boldsymbol{\alpha} \in R_+} k(\boldsymbol{\alpha}). \tag{D15}$$

Using Eq. (D12), one finds the inverse transform of this function to be

$$g_k(\boldsymbol{x}, t) = \frac{1}{c_k} \int_{\mathbb{R}^N} A e^{-t\xi^2/2} E_k(i\boldsymbol{\xi}, \boldsymbol{x}) w_k(\boldsymbol{\xi}) \, \mathrm{d}^N \xi = \frac{A}{t^{\gamma+N/2}} e^{-x^2/(2t)}. \tag{D16}$$

A direct calculation shows that

$$\frac{1}{2} \sum_{j=1}^{N} T_j^2 g_k(\boldsymbol{x}, t) = \frac{g_k(\boldsymbol{x}, t)}{t} \left(\frac{x^2}{t} - N - 2\gamma \right) = \frac{\partial}{\partial t} g_k(\boldsymbol{x}, t). \tag{D17}$$

One could think that the calculation of the Green function is finished by giving g_k a displacement $\boldsymbol{y}$ to obtain

$$g_k(\boldsymbol{y} - \boldsymbol{x}, t) \sim \int_{\mathbb{R}^N} e^{-t\xi^2/2} e^{-i\boldsymbol{\xi}\cdot\boldsymbol{x}} e^{i\boldsymbol{\xi}\cdot\boldsymbol{y}} \, \mathrm{d}^N \xi.$$

However, this function clearly fails to solve Eq. (C2). Instead, we must use the generalized Dunkl translation Eq. (D10),

$$\tau_{-\boldsymbol{x}} g_k(\boldsymbol{y}, t) = \frac{1}{c_k} \int_{\mathbb{R}^N} A e^{-t\xi^2/2} E_k(-i\boldsymbol{x}, \boldsymbol{\xi}) E_k(i\boldsymbol{y}, \boldsymbol{\xi}) w_k(\boldsymbol{\xi}) \,\mathrm{d}^N \xi, \tag{D18}$$

which automatically solves Eq. (C2) on both $\boldsymbol{x}$ and $\boldsymbol{y}$. With the substitution $\boldsymbol{\xi}' = \sqrt{t}\boldsymbol{\xi}$ and using Eq. (D12) we obtain the Green function

$$\Gamma_k(t, \boldsymbol{y}|\boldsymbol{x}) = \tau_{-\boldsymbol{x}} g_k(\boldsymbol{y}, t) = \frac{A e^{-(x^2+y^2)/2t}}{t^{N/2+\gamma}} E_k\left(\frac{\boldsymbol{x}}{\sqrt{t}}, \frac{\boldsymbol{y}}{\sqrt{t}}\right). \tag{D19}$$

In other words, the Dunkl transform of a Gaussian centered at the origin is a Gaussian, but in order to displace the Gaussian correctly and obtain the Green function above, one requires the generalized translation in Eq. (D10).

Note that, since we are interested in calculating a TPD based on Γ_k, it must be normalized. In particular, we must impose the condition that, for an arbitrary starting point $\boldsymbol{x}$ and time t,

$$\int_{\mathbb{R}^N} p_k(t, \boldsymbol{y}|\boldsymbol{x}) \,\mathrm{d}^N y = 1. \tag{D20}$$

As opposed to the case of the N-dimensional Brownian motion, this TPD is not given by Γ_k. For example, if we try to normalize Γ_k directly, we obtain for the case $\boldsymbol{x} = \boldsymbol{0}$

$$1 = \int_{\mathbb{R}^N} \Gamma_k(t, \boldsymbol{y}|\boldsymbol{0}) \,\mathrm{d}^N y = \frac{A}{t^{N/2+\gamma}} \int_{\mathbb{R}^N} e^{-y^2/2t} \,\mathrm{d}^N y = \frac{A(2\pi)^{N/2}}{t^\gamma}. \tag{D21}$$

This leads to $A = t^\gamma/(2\pi)^{N/2}$, which is a clear contradiction because of the presence of t. Therefore, we realize that the normalization of Γ_k must be calculated with respect to w_k in order to obtain a consistent result for A:

$$1 = \int_{\mathbb{R}^N} \Gamma_k(t, \boldsymbol{y}|\boldsymbol{x}) w_k(\boldsymbol{y}) \,\mathrm{d}^N y = A \frac{e^{-x^2/2t}}{t^{N/2+\gamma}} \int_{\mathbb{R}^N} e^{-y^2/2t} E_k\left(\frac{\boldsymbol{x}}{\sqrt{t}}, \frac{\boldsymbol{y}}{\sqrt{t}}\right) w_k(\boldsymbol{y}) \,\mathrm{d}^N y = A c_k. \tag{D22}$$

For this result, we have used Eq. (D12). This yields $A = c_k^{-1}$ and

$$\Gamma_k(t, \boldsymbol{y}|\boldsymbol{x}) = \frac{e^{-(x^2+y^2)/2t}}{c_k t^{\gamma+N/2}} E_k\left(\frac{\boldsymbol{x}}{\sqrt{t}}, \frac{\boldsymbol{y}}{\sqrt{t}}\right). \tag{D23}$$

In view of Eq. (D20), we conclude that

$$p_k(t, \boldsymbol{y}|\boldsymbol{x}) = w_k(\boldsymbol{y}) \Gamma_k(t, \boldsymbol{y}|\boldsymbol{x}) = w_k(\boldsymbol{y}) V_k p_0(t, \boldsymbol{y}|\boldsymbol{x}) = w_k\left(\frac{\boldsymbol{y}}{\sqrt{t}}\right) \frac{e^{-(x^2+y^2)/2t}}{c_k t^{N/2}} E_k\left(\frac{\boldsymbol{x}}{\sqrt{t}}, \frac{\boldsymbol{y}}{\sqrt{t}}\right). \tag{D24}$$

The factor $w_k\left(\boldsymbol{y}/\sqrt{t}\right)$ can be interpreted as follows: in Eq. (C2), the denominators in the second and third terms on the RHS denote a repulsion from the planes defined by $\boldsymbol{\alpha} \cdot \boldsymbol{x} = 0$, $\boldsymbol{\alpha} \in R$, and therefore, the probability of any process finishing at any point such that $\boldsymbol{\alpha} \cdot \boldsymbol{y} = 0$ should be zero.

Indeed, $w_k(\boldsymbol{y}) = \prod_{\boldsymbol{\alpha}\in R} |\boldsymbol{\alpha}\cdot\boldsymbol{y}|^{k(\boldsymbol{\alpha})}$ cancels whenever $\boldsymbol{\alpha}\cdot\boldsymbol{y} = 0$ for any $\boldsymbol{\alpha}\in R$, which reflects this fact.

The existence of V_k is powerful enough to yield all of the above results with relative ease. However, the fact that the explicit form of this operator is unknown imposes a limit to its practical applications.

For the purposes of this work, we are interested in $p_k(t, \boldsymbol{y}|\boldsymbol{x})$ for the root system A. In this case, the Dunkl operator in the direction $\boldsymbol{\xi}$ is given by

$$T_{\boldsymbol{\xi}} f(\boldsymbol{x}) = \partial_{\boldsymbol{\xi}} f(\boldsymbol{x}) + k \sum_{1\leq i<j\leq N} \frac{f(\boldsymbol{x}) - f(\sigma_{ij}\boldsymbol{x})}{x_j - x_i}(\xi_j - \xi_i), \tag{D25}$$

from which Eq. (1) follows. Let us calculate the quantities that form p_k in this particular case. First, the weight function w_k becomes proportional to the Vandermonde determinant $h_N(\boldsymbol{x})$,

$$w_k(\boldsymbol{x}) = \prod_{1\leq j<l\leq N} \left|\frac{1}{\sqrt{2}}(\boldsymbol{e}_l - \boldsymbol{e}_j)\cdot\boldsymbol{x}\right|^{2k} = \frac{|h_N(\boldsymbol{x})|^{2k}}{2^{kN(N-1)/2}}. \tag{D26}$$

The normalization constant c_k is then

$$c_k = \frac{1}{2^{kN(N-1)/2}} \int_{\mathbb{R}^N} e^{-x^2/2} |h_N(\boldsymbol{x})|^{2k} \,\mathrm{d}^N x = \frac{(2\pi)^{N/2}}{2^{kN(N-1)/2}} \prod_{j=1}^{N} \frac{\Gamma(1+jk)}{\Gamma(1+k)}, \tag{D27}$$

where we have used a particular case of the Selberg integral (equation (17.6.7) in Ref. [3]). Inserting the above in Eq. (D24) yields Eq. (28).

Appendix E: Reproduction of Known Results

1. The Intertwining Operator for $k = 0$

From Eqs. (1) and (2), one sees that V_k should reduce to the identity mapping when $k = 0$ in Theorem 2. We prove this in the following proposition.

Proposition 5. *In the limit $k \to 0$, the intertwining operator in Theorem 2 becomes the identity operator, i.e.,*

$$\lim_{k\to 0} V_k m_\lambda(\boldsymbol{x}) = m_\lambda(\boldsymbol{x}). \tag{E1}$$

Proof. Consider the quantity $c_\tau(1/k)/[c'_\tau(1/k)(kN)^{(1/k)}_\tau]$, with $(kN)^{(1/k)}_\tau$ given by Eq. (A16) with $a = kN$ and $\alpha = 1/k$.

$$\frac{c_\tau(1/k)}{c'_\tau(1/k)(kN)^{(1/k)}_\tau} = \prod_{(i,j)\in\tau} \frac{(\tau_i - j + k(\tau'_j - i + 1))}{(k(N-i+1)+j-1)(\tau_i - j + 1 + k(\tau'_j - i))}. \tag{E2}$$

The second factor in the denominator becomes

$$\prod_{(i,j)\in\tau}\frac{1}{\tau_i-j+1+k(\tau'_j-i)}\to\prod_{(i,j)\in\tau}\frac{1}{\tau_i-j+1}=\prod_{i=1}^{l(\tau)}\frac{1}{\tau_i!}=\frac{1}{\tau!} \tag{E3}$$

when k tends to zero. The rest of the expression becomes, after reversing the order of the terms in the numerator,

$$\prod_{i=1}^{l(\tau)}\prod_{j=1}^{\tau_i}\frac{\tau_i-j+k(\tau'_j-i+1)}{k(N-i+1)+j-1}=\prod_{i=1}^{l(\tau)}\prod_{j=1}^{\tau_i}\frac{j-1+k(\tau'_{\tau_i-j+1}-i+1)}{k(N-i+1)+j-1},$$

and after taking the zero limit all terms cancel except for those in which $j=1$:

$$\lim_{k\to 0}\prod_{i=1}^{l(\tau)}\prod_{j=1}^{\tau_i}\frac{j-1+k(\tau'_{\tau_i-j+1}-i+1)}{k(N-i+1)+j-1}=\prod_{i=1}^{l(\tau)}\frac{\tau'_{\tau_i}-i+1}{N-i+1}.$$

The definition of τ'_j is (see Appendix A)

$$\tau'_j=\#\{m|\tau_m\geq j\}$$

for any partition τ (here, # denotes the cardinality of a set), and we are now concerned with

$$\tau'_{\tau_i}=\#\{m|\tau_m\geq\tau_i\}.$$

By the definition of integer partitions, we know that $\tau_m\geq\tau_i$ whenever $m<i$, so τ'_{τ_i} returns the maximum value of m such that $\tau_m=\tau_i$, which we will call l. Here we consider two cases: in the first, $i\neq l$, but then by the definition of partition, i must be less than l and $\tau_i=\tau_{i+1}$. In the second, $i=l$; then, we know immediately that $\tau_i>\tau_{i+1}$. Therefore, for each distinct part in τ, $\tau'_{\tau_i}-i+1$ represents the number of parts τ_m such that $m\geq i$ and $\tau_m=\tau_i$. We then conclude that the product $\prod_{i=1}^{l(\tau)}(\tau'_{\tau_i}-i+1)$ is the product of the factorials of the multiplicities of the non-zero parts of τ. On the other hand, a simple calculation shows that

$$\prod_{i=1}^{l(\tau)}\frac{1}{N-i+1}=\frac{(N-l(\tau))!}{N!},$$

which has the factorial of the multiplicity of the zero parts of τ in the numerator, so we finally arrive at

$$\lim_{k\to 0}\frac{c_\tau(1/k)}{c'_\tau(1/k)(kN)^{(1/k)}_\tau}=\frac{1}{\tau!M(\tau,N)}. \tag{E4}$$

Using the above and the third of Eqs. (A14) the result follows. □

2. The Intertwining Operator for $k = 1$

This case is of interest due to the fact that Jack functions reduce to Schur functions [29], and because of the slightly simplified form that V_k assumes, aside from the fact that it represents the non-colliding Brownian motion ($\beta = 2$).

Proposition 6. *When $k = 1$, the symmetrized Dunkl kernel is given by*

$$\sum_{\rho\in S_N} E_{k=1}(\rho\boldsymbol{x},\boldsymbol{y}) = \prod_{i=1}^{N}(i!)\sum_{\tau:l(\tau)\leq N}\frac{s_\tau(\boldsymbol{x})s_\tau(\boldsymbol{y})}{\prod_{i=1}^{N}(N-i+\tau_i)!} \tag{E5}$$

$$= \prod_{i=1}^{N}(i!)\frac{\det_{1\leq i,j\leq N}(e^{x_i y_j})}{h_N(\boldsymbol{x})h_N(\boldsymbol{y})}. \tag{E6}$$

Furthermore, the TPD of the symmetric Dunkl process is equal to the TPD of the non-colliding Brownian motion [8], Eq. (8)*.*

Proof. At $k = 1$, $u_{\tau\lambda}(1/k)$ becomes the Kostka matrix $K_{\tau\lambda}$, and the functions $c_\tau(1/k)$ and $c'_\tau(1/k)$ become equivalent and cancel in Eq. (31). Also, the generalized Pochhammer symbol, Eq. (A16), becomes

$$(kN)_\tau^{(1/k)}|_{k=1} = \prod_{i=1}^{l(\tau)}\frac{(N-i+\tau_i)!}{(N-i)!}, \tag{E7}$$

which transforms Eq. (31) into [52]

$$V_k m_\lambda(\boldsymbol{x})|_{k=1} = \lambda! M(\lambda,N)\sum_{\substack{\tau:l(\tau)\leq N\\|\tau|=|\lambda|}}\prod_{i=1}^{N}\frac{(N-i)!}{(N-i+\tau_i)!}K_{\tau\lambda}s_\tau(\boldsymbol{x}). \tag{E8}$$

Now, from Eq. (33) and Eq. (E8), we obtain

$$\begin{aligned}\sum_{\rho\in S_N} E_{k=1}(\rho\boldsymbol{x},\boldsymbol{y}) &= \sum_{\mu:l(\mu)\leq N} N! m_\mu(\boldsymbol{y})\sum_{\substack{\tau:l(\tau)\leq N\\|\tau|=|\mu|}}\prod_{i=1}^{N}\frac{(N-i)!}{(N-i+\tau_i)!}K_{\tau\mu}s_\tau(\boldsymbol{x})\\ &= \sum_{\tau:l(\tau)\leq N}\prod_{i=1}^{N}\frac{(N-i)!}{(N-i+\tau_i)!}N! s_\tau(\boldsymbol{x})\sum_{\substack{\mu:l(\mu)\leq N\\|\mu|=|\tau|}}K_{\tau\mu}m_\mu(\boldsymbol{y})\\ &= \sum_{\tau:l(\tau)\leq N}\prod_{i=1}^{N}\frac{(N-i)!}{(N-i+\tau_i)!}N! s_\tau(\boldsymbol{x})s_\tau(\boldsymbol{y}).\end{aligned} \tag{E9}$$

The coefficient of the product of Schur functions can be rewritten as

$$\prod_{i=1}^{N}\frac{(N-i)!}{(N-i+\tau_i)!}N! = \prod_{i=1}^{N}\frac{i!}{(N-i+\tau_i)!}, \tag{E10}$$

from which we obtain Eq. (E5). As shown in Appendix A of Ref. [9], the RHS of Eq. (E5) is the Schur function expansion of the ratio of determinants in Eq. (E6). Therefore, inserting Eq. (E6) in the combination of Eqs. (13) and (28) evaluated at $k = 1$ completes the Karlin-McGregor determinant [53] for the one-dimensional heat kernel and yields Eq. (8). $\square$

3. Dyson's Model: Particles Starting From the Same Position

In view of the A-type Dunkl operator, Eq. (D25), we see that when it is calculated along the vector $\boldsymbol{\xi} = \mathbf{1}$, the difference term cancels and

$$T_{\mathbf{1}} f(\boldsymbol{x}) = \partial_{\mathbf{1}} f(\boldsymbol{x}) \tag{E11}$$

for an arbitrary differentiable function f. From this, one can deduce that the intertwining operator should have no effect on f along the line defined by $\mathbf{1}$. Moreover, the Dunkl kernel should reduce to a regular exponential in this case. The consequence of this fact on Theorem 2 is the following.

Proposition 7. *The intertwining operator has no effect on the monomial symmetric function* $m_\lambda(\boldsymbol{x})$ *when it is evaluated at* $\boldsymbol{x} = x_0\mathbf{1}$, *where* $x_0 \in \mathbb{R}$. *Moreover, the TPD of the symmetric Dunkl process whose initial configuration is* $\boldsymbol{x} = x_0\mathbf{1}$ *is given by*

$$p_k^s(t, \boldsymbol{y}|x_0\mathbf{1}) = p_k^s(t, \boldsymbol{y} - x_0\mathbf{1}|\mathbf{0}) = \frac{N!}{(2\pi t)^{N/2}} \prod_{j=1}^{N} \frac{\Gamma(1+k)}{\Gamma(1+jk)} |h_N(\boldsymbol{y})|^{2k} e^{-(\boldsymbol{y}-x_0\mathbf{1})^2/2t}. \tag{E12}$$

Proof. First, we prove that $V_k m_\lambda(\boldsymbol{x})$ is independent of k whenever $\boldsymbol{x}$ is a multiple of $\mathbf{1}$. For this purpose, we derive an expression which will prove useful for our calculations. By definition, the C-normalized Jack functions satisfy Eq. (A10), which is equivalent to

$$\sum_{\substack{\tau: l(\tau)\leq N\\ |\tau|=n}} \mathcal{C}_\tau^{(\alpha)}(\boldsymbol{x}) = (x_1 + x_2 + \ldots + x_N)^n \tag{E13}$$

for any non-negative integer n. The RHS can be rewritten in terms of monomial symmetric functions,

$$(x_1 + x_2 + \ldots + x_N)^n = \sum_{\substack{\nu: l(\nu)\leq N\\ |\nu|=n}} \frac{n!}{\nu!} m_\nu(\boldsymbol{x}), \tag{E14}$$

but at the same time, we can rewrite the LHS of Eq. (E13) using Eqs. (A18) and (A11):

$$\sum_{\substack{\tau: l(\tau)\leq N\\ |\tau|=n}} \mathcal{C}_\tau^{(\alpha)}(\boldsymbol{x}) = \sum_{\substack{\tau: l(\tau)\leq N\\ |\tau|=n}} \frac{|\tau|!}{k^{|\tau|} c'_\tau(1/k)} \mathcal{P}_\tau^{(1/k)}(\boldsymbol{x}) = \sum_{\substack{\tau: l(\tau)\leq N\\ |\tau|=n}} \frac{n!}{k^n c'_\tau(1/k)} \sum_{\substack{\mu: l(\mu)\leq N\\ |\mu|=n}} u_{\tau\mu}(\alpha) m_\mu(\boldsymbol{x}). \tag{E15}$$

Since the monomial symmetric functions are linearly independent, we can equate the coefficients of m_ν in Eqs. (E14) and (E15) to obtain

$$\sum_{\substack{\tau:l(\tau)\leq N\\|\tau|=n}} \frac{u_{\tau\nu}(\alpha)}{c'_\tau(1/k)} = \frac{k^n}{\nu!}. \tag{E16}$$

Now we consider Theorem 2 at $\boldsymbol{x} = x_0\mathbf{1}$.

$$\begin{aligned} V_k m_\lambda(x_0\mathbf{1}) &= \lambda! M(\lambda, N) \sum_{\substack{\tau:l(\tau)\leq N\\|\tau|=|\lambda|}} \frac{u_{\tau\lambda}(1/k)}{c'_\tau(1/k)} \left(\frac{x_0}{k}\right)^{|\tau|} \\ &= \lambda! M(\lambda, N) \frac{x_0^{|\lambda|}}{k^{|\lambda|}} \sum_{\substack{\tau:l(\tau)\leq N\\|\tau|=|\lambda|}} \frac{u_{\tau\lambda}(1/k)}{c'_\tau(1/k)}. \end{aligned} \tag{E17}$$

We insert Eq. (E16) with $n = |\lambda|$ and $\nu = \lambda$ to obtain

$$V_k m_\lambda(x_0\mathbf{1}) = M(\lambda, N) x_0^{|\lambda|}. \tag{E18}$$

As expected, this quantity is independent of k. Note that this is the same value as

$$m_\lambda(x_0\mathbf{1}) = x_0^{|\lambda|} \sum_{\substack{\text{distinct}\\\text{permutations of } \lambda}} 1 = M(\lambda, N) x_0^{|\lambda|}, \tag{E19}$$

which shows that V_k has no effect on m_λ along $\mathbf{1}$. Therefore,

$$\sum_{\rho\in S_N} E_k(\rho(x_0\mathbf{1}), \boldsymbol{y}) = N! e^{x_0\mathbf{1}\cdot\boldsymbol{y}}. \tag{E20}$$

Inserting the above in Eq. (13) combined with Eq. (28) evaluated at $\boldsymbol{x} = x_0\mathbf{1}$ completes the proof. □

This is consistent with previous results [4, 7]. In particular, when $x_0 = 0$ and $t = 1/(2k)$, the TPD in Eq. (E12) becomes

$$p_k^s(1/(2k), \boldsymbol{y}|\mathbf{0}) = N! \frac{2^{\frac{1}{2}kN(N-1)} k^{\frac{1}{2}N+\frac{1}{2}kN(N-1)}}{\pi^{N/2}} \prod_{j=1}^{N} \frac{\Gamma(1+k)}{\Gamma(1+jk)} \left|h_n(\boldsymbol{y})\right|^{2k} e^{-ky^2}. \tag{E21}$$

This is the joint probability density function of the eigenvalues $\boldsymbol{y}$ of a matrix drawn from the $\beta = 1, 2, 4$ ensembles of Gaussian random matrices multiplied by a factor of $N!$, with $\beta = 2k$ [3].

Appendix F: Extema and Maximum value of $F_N(\boldsymbol{v}, t)$

Here, we prove two lemmas necessary to complete the proof of Theorem 4. The first concerns the location of the maxima of $F_N(\boldsymbol{v}, t)$.

Lemma 8. *The maxima of the function*

$$F_N(\boldsymbol{v},t) = \frac{N}{2}(N-1)(1-\log t) - \sum_{j=1}^{N} j\log j + 2\log|h_N(\boldsymbol{v})| - \frac{v^2}{2t} \tag{F1}$$

are located at $\boldsymbol{v} = \sqrt{2t}\boldsymbol{z}_N$ *or any of its permutations, where* $\boldsymbol{z}_N$ *is the vector of roots of the Hermite polynomial* $H_N(x)$*, given by Eqs.* (51) *and* (50) *respectively.*

Proof. We first calculate the first-order partial derivatives of $F_N(\boldsymbol{v},t)$ relative to $\boldsymbol{v}$ and equate them to zero.

$$\frac{\partial}{\partial v_i} F_N(\boldsymbol{v},t) = \sum_{\substack{j:j\neq i\\ j=1}}^{N} \frac{2}{v_i - v_j} - \frac{v_i}{t} = 0 \tag{F2}$$

Hence, the extrema of $F_N(\boldsymbol{v},t)$ must obey the relation

$$v_i = \sum_{\substack{j:j\neq i\\ j=1}}^{N} \frac{2t}{v_i - v_j}. \tag{F3}$$

Furthermore, applying the scaling $\boldsymbol{v} = \sqrt{2t}\boldsymbol{z}$ we obtain

$$z_i = \sum_{\substack{j:j\neq i\\ j=1}}^{N} \frac{1}{z_i - z_j}. \tag{F4}$$

Therefore, it suffices to solve the above equation for $\boldsymbol{z}$ to find the location of the extrema of $F_N(\boldsymbol{v},t)$. The second order derivatives of $F_N(\boldsymbol{v},t)$ are

$$\frac{\partial^2}{\partial v_j \partial v_i}\left[2\log|h_N(\boldsymbol{v})| - \frac{v^2}{2t}\right] = \frac{\partial}{\partial v_j}\left[\sum_{l:l\neq i} \frac{2}{v_i - v_l} - \frac{v_i}{t}\right] = \begin{cases} -\sum_{l:l\neq i} \frac{2}{(v_i-v_l)^2} - \frac{1}{t} & \text{if } i = j, \\ \frac{2}{(v_i-v_j)^2} & \text{if } i \neq j. \end{cases} \tag{F5}$$

The matrix formed by the $N\times N$ second order derivatives above is negative definite for all vectors $\boldsymbol{v}$ with non-repeating components. To show this, we consider an arbitrary real vector $\boldsymbol{u}$ and calculate the quadratic form associated to Eq. (F5).

$$\sum_{1\leq i,j\leq N} u_i \frac{\partial^2 F_N(\boldsymbol{v},t)}{\partial v_j \partial v_i} u_j = -\frac{1}{t}\sum_{i=1}^{N} u_i^2 - 2\sum_{1\leq i<j\leq N} \frac{(u_i-u_j)^2}{(v_i-v_j)^2} \leq 0 \tag{F6}$$

Here, the equality holds only when all the u_i are equal to zero. This shows that all extrema given by Eq. (F3) are maxima.

Note that, given $\boldsymbol{z}$, any of its permutations solve Eq. (F4):

$$z_{\rho(i)} = \sum_{\substack{j:j\neq i\\ j=1}}^{N} \frac{1}{z_{\rho(i)} - z_{\rho(j)}} \tag{F7}$$

for any $\rho \in S_N$. Now, we prove that the roots of the Hermite polynomial H_N solve Eq. (F4). One can write H_N as follows:

$$H_N(x) = 2^N \prod_{j=1}^{N} (x - z_{j,N}). \tag{F8}$$

The derivative of the logarithm of H_N is

$$\frac{H_N'(x)}{H_N(x)} = \sum_{j=1}^{N} \frac{1}{x - z_{j,N}}, \tag{F9}$$

where

$$H_N'(x) = \frac{\mathrm{d}}{\mathrm{d}x} H_N(x). \tag{F10}$$

We subtract the i-th term of the sum from both sides to obtain

$$\frac{H_N'(x)}{H_N(x)} - \frac{1}{x - z_{i,N}} = \sum_{\substack{j:j=1 \\ j\neq i}}^{N} \frac{1}{x - z_{j,N}}. \tag{F11}$$

Taking the limit $x \to z_{i,N}$ yields

$$\sum_{\substack{j:j\neq i \\ j=1}}^{N} \frac{1}{z_{i,N} - z_{j,N}} = \lim_{x \to z_{i,N}} \frac{H_N'(x)}{H_N(x)} - \frac{1}{x - z_{i,N}}. \tag{F12}$$

At the same time,

$$H_N'(z_{i,N}) = \lim_{x \to z_{i,N}} \frac{H_N(x)}{x - z_{i,N}} = 2^N \prod_{\substack{j:j\neq i \\ j=1}}^{N} (z_{i,N} - z_{j,N}). \tag{F13}$$

We will denote the expression in the limit by

$$H_{i,N}(x) = 2^N \prod_{\substack{j:j\neq i \\ j=1}}^{N} (x - z_{j,N}) = \frac{H_N(x)}{x - z_{i,N}} \tag{F14}$$

so that

$$\lim_{x \to z_{i,N}} H_{i,N}(x) = H_N'(z_{i,N}). \tag{F15}$$

With this, Eq. (F12) can be rewritten as

$$\begin{aligned} \sum_{\substack{j:j\neq i \\ j=1}}^{N} \frac{1}{z_{i,N} - z_{j,N}} &= \lim_{x \to z_{i,N}} \frac{1}{H_{i,N}(x)} \left(\frac{H_N'(x)}{x - z_{i,N}} - \frac{H_{i,N}(x)}{x - z_{i,N}} \right) \\ &= \lim_{x \to z_{i,N}} \frac{1}{H_{i,N}(x)} \left(\frac{H_N'(x)}{x - z_{i,N}} - \frac{H_N(x)}{(x - z_{i,N})^2} \right). \end{aligned} \tag{F16}$$

In the vicinity of $z_{i,N}$, $H_N(x)$ is approximately equal to

$$H_N(x) \approx H_N'(z_{i,N})(x - z_{i,N}) + \frac{1}{2}H_N''(z_{i,N})(x - z_{i,N})^2, \tag{F17}$$

where

$$H_N''(x) = \frac{\mathrm{d}^2}{\mathrm{d}x^2}H_N(x). \tag{F18}$$

At the same time, Hermite polynomials satisfy the differential equation [46]

$$H_N''(x) = 2xH_N'(x) - 2NH_N(x), \tag{F19}$$

which implies that $H_N''(z_{i,N}) = 2z_{i,N}H_N'(z_{i,N})$, so

$$H_N(x) \approx H_N'(z_{i,N})(x - z_{i,N})[1 + z_{i,N}(x - z_{i,N})]. \tag{F20}$$

Inserting the above in Eq. (F16) yields

$$\begin{aligned}\sum_{\substack{j:j\neq i\\ j=1}}^{N} \frac{1}{z_{i,N} - z_{j,N}} &= \lim_{x\to z_{i,N}} \frac{1}{H_{i,N}(x)}\left(\frac{H_N'(x) - H_N'(z_{i,N})}{x - z_{i,N}} - z_{i,N}H_N'(z_{i,N})\right)\\ &= \frac{H_N''(z_{i,N})}{H_N'(z_{i,N})} - z_{i,N} = z_{i,N},\end{aligned} \tag{F21}$$

which completes the proof. □

In the second lemma, we find the maximum value of $F_N(\boldsymbol{v}, t)$.

Lemma 9. *The maximum value of* $F_N(\boldsymbol{v}, t)$ *is zero.*

Proof. By Lemma 8, the maximum value of $F_N(\boldsymbol{v}, t)$ is located at $\boldsymbol{v} = \sqrt{2t}\boldsymbol{z}_N$. This yields

$$F_N(\sqrt{2t}\boldsymbol{z}_N, t) = \frac{N}{2}(N-1)(1 + \log 2) - \sum_{j=1}^{N} j\log j + 2\log|h_N(\boldsymbol{z}_N)| - z_N^2, \tag{F22}$$

which is independent of t. Let us focus first on the term

$$z_N^2 = \sum_{j=1}^{N} z_{j,N}^2. \tag{F23}$$

The derivative property of the Hermite polynomials [46] allows us to write

$$H_N'(x) = 2NH_{N-1}(x), \tag{F24}$$

and using the product representation from Lemma 8 we obtain

$$\sum_{i=1}^{N} \prod_{\substack{j:j=1\\ j\neq i}}^{N} (x - z_{j,N}) = N\prod_{j=1}^{N-1}(x - z_{j,N-1}). \tag{F25}$$

Since the above equation holds for all x, we can expand in powers of x and equate the coefficients of the same order. In particular, the coefficients of x^{N-3} form the equation

$$(N-2)\sum_{1\le i<j\le N} z_{i,N}z_{j,N} = N\sum_{1\le i<j\le N-1} z_{i,N-1}z_{j,N-1}, \tag{F26}$$

and if we denote the double sum on the LHS by s_N, we obtain

$$s_N = \frac{N}{N-2}s_{N-1}. \tag{F27}$$

Now, we can show that $s_N = -N(N-1)/4$ by induction. Since $H_2(x) = 4x^2 - 2$, its two roots lie at $x = \pm\sqrt{1/2}$ and therefore $s_2 = -1/2$. Then, the formula is valid for $N = 2$. Assuming it is also valid for $N-1$, we verify that it holds for N:

$$s_N = \frac{N}{N-2}s_{N-1} = -\frac{N}{N-2}\frac{(N-1)(N-2)}{4} = -\frac{N(N-1)}{4}, \tag{F28}$$

as desired. Now, since the Hermite polynomials have either odd or even symmetry, the sum of their roots equals zero. Therefore,

$$0 = \left(\sum_{i=1}^{N} z_{i,N}\right)^2 = z_N^2 + 2s_N = z_N^2 - 2\frac{N(N-1)}{4}, \tag{F29}$$

which finally yields

$$z_N^2 = \frac{N(N-1)}{2}. \tag{F30}$$

Now, let us calculate

$$2\log|h_N(\boldsymbol{z}_N)|. \tag{F31}$$

The main quantity we wish to find is the square of the Vandermonde determinant of the roots of the N-th Hermite polynomial. This quantity is known as the discriminant of the Hermite polynomials, and we calculate it here following Szegö [47]. Using Eq. (F13), we can write

$$\begin{aligned}(h_N(\boldsymbol{z}_N))^2 &= \prod_{1\le i<j\le N}(z_{j,N}-z_{i,N})^2 = (-1)^{N(N-1)/2}\prod_{1\le i\ne j\le N}(z_{j,N}-z_{i,N}) \\ &= \frac{(-1)^{N(N-1)/2}}{2^{N^2}}\prod_{i=1}^{N}H_N'(z_{i,N}) = \frac{(-1)^{N(N-1)/2}N^N}{2^{N(N-1)}}\prod_{i=1}^{N}H_{N-1}(z_{i,N}).\end{aligned} \tag{F32}$$

The last equality follows from Eq. (F24). Let us focus on the last product:

$$\begin{aligned}\prod_{i=1}^{N}H_{N-1}(z_{i,N}) &= 2^{N(N-1)}\prod_{i=1}^{N}\prod_{j=1}^{N-1}(z_{i,N}-z_{j,N-1}) = 2^{N(N-1)}\prod_{j=1}^{N-1}\prod_{i=1}^{N}(z_{j,N-1}-z_{i,N}) \\ &= \prod_{j=1}^{N-1}H_N(z_{j,N-1}).\end{aligned} \tag{F33}$$

From the recurrence relation

$$H_N(x) = 2xH_{N-1}(x) - 2(N-1)H_{N-2}(x) \tag{F34}$$

we obtain that $H_N(z_{j,N-1}) = -2(N-1)H_{N-2}(z_{j,N-1})$, so the product above becomes

$$\prod_{i=1}^{N} H_{N-1}(z_{i,N}) = [-2(N-1)]^{N-1} \prod_{j=1}^{N-1} H_{N-2}(z_{j,N-1}). \tag{F35}$$

For $N = 2$, the LHS is equal to

$$H_1(-1/\sqrt{2})H_1(1/\sqrt{2}) = -2, \tag{F36}$$

for which the general formula

$$\prod_{i=1}^{N} H_{N-1}(z_{i,N}) = (-2)^{N(N-1)/2} \prod_{j=1}^{N-1} j^j \tag{F37}$$

holds. Now, assuming that Eq. (F37) holds for $N-1$, we verify that it holds for N:

$$\begin{aligned}\prod_{i=1}^{N} H_{N-1}(z_{i,N}) &= [-2(N-1)]^{N-1} \prod_{j=1}^{N-1} H_{N-2}(z_{j,N-1}) \\ &= [-2(N-1)]^{N-1}(-2)^{(N-1)(N-2)/2} \prod_{j=1}^{N-2} j^j = (-2)^{N(N-1)/2} \prod_{j=1}^{N-1} j^j.\end{aligned} \tag{F38}$$

Therefore,

$$(h_N(\boldsymbol{z}_N))^2 = \frac{(-1)^{N(N-1)/2} N^N}{2^{N(N-1)}} \prod_{i=1}^{N} H_{N-1}(z_{i,N}) = \frac{1}{2^{N(N-1)/2}} \prod_{j=1}^{N} j^j. \tag{F39}$$

The logarithm of the above is

$$2\log|h_N(\boldsymbol{z}_N)| = \sum_{j=1}^{N} j\log j - \frac{N}{2}(N-1)\log 2, \tag{F40}$$

and thus the maximum value of $F_N(\boldsymbol{v}, t)$ is zero for all its extrema, due to the fact that Eqs. (F23) and (F31) do not change when the roots $\boldsymbol{z}_N$ are permuted. □

---

[1] F. J. Dyson, J. Math. Phys. **3**, 1191 (1962).

[2] F. J. Dyson, J. Math. Phys. **3**, 1199 (1962).

[3] M. L. Mehta, *Random Matrices* (Elsevier, 2004).

[4] P. J. Forrester, *Log-Gases and Random Matrices* (Princeton University Press, 2010).

[5] M. Katori and H. Tanemura, J. Stat. Phys. **129**, 1233 (2007).

[6] M. E. Fisher, J. Stat. Phys. **34**, 667 (1984).

[7] M. Katori and H. Tanemura, Phys. Rev. E **66** (2002).

[8] D. J. Grabiner, Ann. I.H.P. (B), Prob. Stat. **35**, 177 (1999).

[9] M. Katori and H. Tanemura, J. Math. Phys. **45**, 3058 (2004).

[10] J. W. Essam and A. J. Guttmann, Phys. Rev. E **52**, 5849 (1995).

[11] P. de Gennes, J. Chem. Phys. **48**, 2257 (1968).

[12] J. Baik, A. Borodin, P. Deift, and T. Suidan, J. Phys. A: Math. Gen. **39**, 8965 (2006).

[13] O. Bohigas, R. Haq, and A. Pandey, in *Nuclear Data for Science and Technology*, edited by K. Böckhoff (D. Reidel, 1983).

[14] O. Bohigas, R. U. Haq, and A. Pandey, Phys. Rev. Lett. **54**, 1645 (1985).

[15] S. de Haro, Nucl. Phys. B **730**, 312 (2005).

[16] P. J. Forrester, S. N. Majumdar, and G. Schehr, Nucl. Phys. B. **844**, 500 (2011).

[17] S. de Haro and M. Tierz, Phys. Lett. B **601**, 201 (2004).

[18] K. Johansson, Comm. Math. Phys. **209**, 437 (2000), 10.1007/s002200050027.

[19] K. Johansson, Comm. Math. Phys. **242**, 277 (2003).

[20] M. Prähofer and H. Spohn, Phys. Rev. Lett. **84**, 4882 (2000).

[21] T. Imamura and T. Sasamoto, Phys. Rev. E **71**, 041606 (2005).

[22] T. Sasamoto and H. Spohn, Phys. Rev. Lett. **104**, 230602 (2010).

[23] T. Imamura and T. Sasamoto, J. Phys. A: Math. Theor. **44**, 385001 (2011).

[24] K. A. Takeuchi and M. Sano, Phys. Rev. Lett. **104**, 230601 (2010).

[25] K. A. Takeuchi, M. Sano, T. Sasamoto, and H. Spohn, Sci. Rep. **1** (2011).

[26] A. J. Guttmann, A. L. Owczarek, and X. G. Viennot, J. Phys. A: Math. Gen. **31**, 8123 (1998).

[27] C. Krattenthaler, A. J. Guttmann, and X. G. Viennot, J. Phys. A: Math. Gen. **33**, 8835 (2000).

[28] W. Fulton, *Young tableaux: with applications to representation theory and geometry* (Cambridge University Press, 1997).

[29] I. G. Macdonald, *Symmetric Functions and Hall Polynomials* (Oxford University Press, 1995).

[30] C. F. Dunkl, Trans. Amer. Math. Soc. **311**, 167 (1989).

[31] C. F. Dunkl and Y. Xu, *Orthogonal Polynomials of Several Variables* (Cambridge University Press, 2001).

[32] K. Hikami, J. Phys. Soc. Jpn. **65**, 394 (1996).

[33] L. Lapointe and L. Vinet, Comm. Math. Phys. **178**, 425 (1996).

[34] C. F. Dunkl, J. Phys. A: Math. Gen. **35**, 10391 (2002).

[35] C. F. Dunkl, in *Hypergeometric functions on domains of positivity, Jack polynomials, and applications (Tampa, FL, 1991)*, Contemp. Math., Vol. 138 (Amer. Math. Soc., Providence, RI, 1992) pp. 123–138.

[36] M. Rösler, Comm. Math. Phys. **192**, 519 (1998).

[37] M. Rösler and M. Voit, in *Harmonic & stochastic analysis of Dunkl processes*, edited by P. Graczyk, M. Rösler, and M. Yor (HERMANN Mathématiques, 2008).

[38] N. Demni, in *Harmonic & stochastic analysis of Dunkl processes*, edited by P. Graczyk, M. Rösler, and M. Yor (HERMANN Mathématiques, 2008).

[39] N. Demni, SIGMA **4**, 74 (2008).

[40] N. Demni, C. R. Math. **347**, 1125 (2009).

[41] N. Demni, in *Séminaire de Probabilités XLII*, edited by C. Donati-Martin, M. Émery, A. Rouault, and C. Stricker (Springer, 2009).

[42] C. F. Dunkl, Can. J. Math. **43**, 1213 (1991).

[43] T. Baker and P. Forrester, Comm. Math. Phys. **188**, 175 (1997).

[44] E. M. Opdam, Compo. Math. **85**, 333 (1993).

[45] M. Rösler, Duke Math. J. **98**, 445 (1999).

[46] G. Arfken and H. Weber, *Mathematical Methods for Physicists* (Elsevier, 2005).

[47] G. Szegö, *Orthogonal Polynomials* (American Mathematical Society, 1975).

[48] W. Feller, *An Introduction to Probability Theory and Its Applications* (Wiley, 1968).

[49] N. Demni, SIGMA **4**, 75 (2008).

[50] R. P. Stanley, Adv. Math. **77**, 76 (1989).

[51] I. Dumitriu, A. Edelman, and G. Shuman, J. Symb. Comp. **42**, 587 (2007).

[52] S. Andraus, *Dunkl Processes and Intertwining Operators*, Master's thesis, Chuo University (2011).

[53] S. Karlin and J. McGregor, Pac. J. Math **9**, 1141 (1959).